


ABSTRACTED IN CABI, U.K., SHIMAGO, CROSS REF,
INDIAN CITATION INDEX AND GOOGLE SCHOLAR,
PUBLONS, CITEFACTOR



Journal of Natural Resource And Development

(Peer Reviewed, Refereed Research Journal of Agriculture and Science)

Abbreviated title of Journal : *Jour. Nat. Res. Dev.*

© copyright. Editor, SBSRD, Prayagraj, (U.P.), India

NAAS RATING : 3.77

online @ www.sbsrd.org

SOCIETY OF BIOLOGICAL SCIENCES AND RURAL DEVELOPMENT

10/96, Gola Bazar, New Jhusi, Prayagraj - 211 019 (U.P.), INDIA

EXECUTIVE COUNCIL

Patron Dr. S. C. Pathak	President Prof. Krishna Mishra	Secretary Dr. Hemlata Pant
-----------------------------------	--	--------------------------------------

ADVISORY BOARD

- Prof. Shyam Narayan Labh**, Professor & Head
Campus Research Committee (CRC), B Amrit Campus, Tribhuvan University,
Kathmandu - 44600, Nepal
- Dr. B. K. Chakraborty**
Former Director, Department of Fisheries, Bangladesh and Supervisor/Advisor,
Bangladesh Agricultural University, Bangladesh
- Dr. J.C. Tarafdar**
Ex-Principal Scientist, ICAR-CAZRI, Jodhpur-342 003 (Rajasthan), India
- Dr. Y. P. Singh**
Principal Scientist & Head, ICAR-Central Soil Salinity Research Institute,
Dilkusha, Lucknow 226002 (UP) India
- Dr. A.S. Ninawe**, Ex-Senior Advisor,
Department of Biotechnology, New Delhi, India
- Dr. P. Keshav Nath**
Former Deen, Fisheries Karnataka Veterinary, Animal & Fisheries Sciences, Bidar, India
- Dr. Eduardo Lobo Alcayaga**,
Department of Biology and Pharmacy, UNISC, Brazil
- Dr. D.V. Singh**, Professor and Head,
LPM, GBPUAT, Pantnagar, Uttrakhand, India
- Prof. Krishna Kumar**, Ex Dean Science,
University of Allahabad, Prayagraj, (U.P.), India
- Prof. Prakash Nautiyal**, Department of Zoology and Biotechnology,
HNB Garhwal University, Srinagar, (U.K.), India
- Dr. A. Arunachalam**, Director, Central Agroforestry Research Institute (ICAR), Jhansi, (U.P.), India
- Prof. A.R. Siddiqui**, Head, Department of Geography,
University of Allahabad, Prayagraj, (U.P.), India
- Prof. Vipin Masih Prasad**
Prof., Ex-Head, Department of Horticulture, SHUATS, Prayagraj, U.P., India

EDITORIAL BOARD

Chief-Editor : Dr. Hemlata Pant Prayagraj, (U.P.), India	Editors : Prof. Surya Narayan , Prayagraj, (U.P.), India Dr. A.S. Yadav , Kanpur Nagar, (U.P.), India Dr. Manoj Kumar Singh , Prayagraj, (U.P.), India Dr. Sandeep Kushwaha , Jabalpur, (M.P.), India Dr. Ashwani Kumar , Mathura, (M.P.), India	Associate Editor : Dr. Jyoti Verma Prayagraj, (U.P.), India
--	--	---

EDITORIAL BOARD MEMBERS

Dr. Ramesh D. Gulati

Senior Emeritus Scientist, Netherlands Institute of Ecology,
Department of Aquatic Ecology, Netherlands

Dr. U. K. Sarkar

Director, ICAR - NBFGR, Dilkusha Road, Lucknow, (U.P.), India

Dr. D. Prasad

Ex-Principal, Scientist and Head, Division of Nematology, IARI, New Delhi, India

Dr. A.K. Pandey

Dean, College of Horticulture, Central Agriculture University, Bermiok, South Sikkim - 737134, India

Dr. Narendra Kumar

Principal Scientist (Agronomy), Division of Crop Production,
ICAR-Indian Institute of Pulses Research, Kanpur- 208 024 (U.P.), India,

Dr. Ranjit Singh Yadav

Principal Scientist (Soil Science), ICAR-CAZRI, Jodhpur-342 003 (Rajasthan), India

Dr. Sanjay Arora

Principal Scientist (Soil Science), ICAR-Central Soil Salinity Research Institute,
Jail Road, Lucknow 226002 (UP) India

Prof, K.P. Singh

Head, Dept. of Zoology, University of Allahabad, Prayagraj, (U.P.), India

Dr. D.K. Srivastava

Joint Director, Agriculture, CSTUP, Lucknow, (U.P.), India

Dr. Manish Kanwat

Principal Scientist, ICAR - Central Arid Zone Research Institute, Kutch (Gujrat), India

Dr. Rakesh Kumar Dubey

Principal Scientist, ICAR - Indian Institute of Vegetable Research, Varanasi, (U.P.), India

Dr. Deepika Slathia

Department of Environmental Sciences, University of Jammu, Jammu-180006. (J&K), India

Dr. Deepak Kumar Srivastava

Principal, Carreer Convent Girl P.G. College, Lucknow, U.P., India

Dr. K. Dinesh

Asso. Prof. and Head, Fisheries Station,
Kerala University of Fisheries and Ocean Studies (KUFOS), Kochi, Kerala, India

Dr. R.K. Naik

Senior Scientist, Farm Machinery and Power Agri. Engg. Section,
(ICAR)-Central Research Institute for Jute and allied Fibres,
Nilganj, Barrackpore, Kolkata, (W.B.)

Prof. Harikesh Singh

Principal, Ganna Utpadak Mahavidyalaya, Baheri, Bareilly, (U.P.), India

Prof. S.N. Sharma

Principal, Meerut College, Meerut, (U.P.), India

Prof. Laxmi Yadav

Department of Zoology, KPM Govt. Womens Degree College, Aurai, Bhadohi (U.P.), India

Dr. D.K. Chauhan

Asso. Prof., Department of Zoology, CCS University, Meerut, (U.P.), India

Dr. Ashok Kumar Singh

Asso. Prof., Dept. of Plant Pathology, SKUAST - J, Chatha, Jammu, India

Dr. S.P. Singh

Asso.Prof., Agricultural Economics & ABM, SKUAST, Jammu, India

Dr. Neeraj Gupta

Asso. Prof., Dept. of Post Harvest Technology, SKUAST Jammu, India

Dr. S.P. Verma

Prof., Dept. of A. H. & D., KAPG College, Prayagraj, (U.P.), India

Dr. D. Swaroop

Animal Scientist/Asst. Prof. CSAUA & T, Kanpur, (U.P.), India

Dr Maheshwaree Prasad Singh

Incharge KVK Chhata Prayagraj II

Dr. Rakesh Dwivedi

Asst. Prof., Department of Botany, Higher Education, Jaihari Khal (U.K.)

Dr. Jitendra Kumar Shukla

Asst. Director Fisheries, Bahraich, (Govt. of U.P.), U.P., India

Dr. Dharmendra Singh

Asst. Prof., Dept. of Zoology, Goswami Tulsidad Govt. P.G. College, Karwi, Chitrakoot, (U.P.), India

Dr. Pramod Kumar

Asst. Prof., Dept. of Chemistry, Government Degree College, Manikpur Chitrakoot, (U.P.), India

Dr. Varsha Jaiswal

Asst. Prof., Dept. of Botany, PBPG College, Pratapgarh, (U.P.), India

Dr. Anita Singh

Dept. of Botany, CMP PG College, Prayagraj, (U.P.), India

Dr. Balaji Vikram

Asst. Prof., Dept. of Post Harvest Technology, College of Horticulture,
Banda University of Agriculture and Technology, Banda, U.P., India-210001

REVIEWER COMMITTEE

Dr. Hemant Kumar

Senior Scientist, ICAR--IIPR, Kanpur, (U.P.), India

Dr. A.K. Singh

Asst. Prof., Dept. Entomology, Buant, Banda, (U.P.), India

Dr. O.P. Maurya

Asso. Prof., Dept. of Agricultural Economics,
RSM P.G. College, Dhampur, Bijnor, (U.P.), India

Dr. Kirti Raje Singh

Asst. Prof., Dept. of Botany, Prayagraj, (U.P.), India

Dr. Pallavi Rai

Asst. Prof., Dept. of Botany, CMP PG College, Prayagraj, (U.P.), India

Dr. Archana Rai

NIET, NIMS University, Rajasthan Campus, Jaipur, Rajasthan, India

Note : The above members are not salaried from this organization

SOCIETY OF BIOLOGICAL SCIENCES AND RURAL DEVELOPMENT

CONTENTS

➡	A COMPARISON OF OCCURRENCE OF FOSSILS OF P.S. Bhatnagar, and S. Pujari	1-5
➡	INTER CROPPING OF SUGARCANE AND CAULIFLOWER Pradeep Kumar Misra, Manoj Kumar Singh and R L Yadav	6-10
➡	AN ECO-FRIENDLY APPROACH TO BIOREMEDIATION Barkha Kamal, Sujata Gupta, Reena Verma, M. M. Jwantha and M. K. Purohit	11-16
➡	NUTRITIONAL IMPORTANCE AND PROCESSING ASPECTS Niraj Kumar, Neeraj Singh and Vivek Singh	17-23
➡	BASELINE DATA ON BIRD DIVERSITY IN PRAYAGRAJ Imran Alam, Md. Noor Alam and Sandeep Kushwaha	24-29
➡	EXPLORING CONSTRAINTS FACED BY FARMERS IN THE Anish Kumar and N. K. Mishra	30-34
➡	IMPACT OF DIFFERENT PHOSPHATE SUBSTRATES Meenakshee Pandey, Shalini Singh and Shweta Mishra	35-44
➡	ASSESSING THE CHALLENGES HINDERING FARM WOMEN'S Km. Arti, N. K. Mishra, and Pradeep Kumar Yadav	45-48
➡	IMPACT OF KVK TRAINING PROGRAM ON KNOWLEDGE Anish Kumar, N. K. Mishra and Pradeep Kumar Yadav	49-53
➡	EFFECT OF DIFFERENT VARIABLES ON PHYSICO-CHEMICAL Siddhartha Singh, Akhilesh Kumar Singh, Manoj Kumar, Manvendra Singh, Rajesh Kumar Pal and Kautuk Upadhyay	54-60
➡	MOTHS (LEPIDOPTERA: HETEROCERA) DIVERSITY OF S. Sambath and Akhil Nair	61-69
➡	PREVALENCE OF MALNUTRITION AMONG PRESCHOOL Shriya Singh, N. K. Mishra and Pradeep Kumar Yadav	70-73
➡	USE OF BOTANICAL PESTICIDES IN SUSTAINABLE Preeti Maurya and Sanjay Singh	74-85
➡	EFFECT OF INTEGRATED NUTRIENT MANAGEMENT Manoj Kumar Singh	86-89
➡	PRODUCTIVE, REPRODUCTIVE STATUS NUTRIENT AVAILABILITY Vidya Sagar, Pradeep Kumar, Ram Jeet and Ram Gopal Yadava	90-95
➡	CONTENT ANALYSIS OF AGRICULTURE INFORMATION Km. Bhawana Singh and N. K. Mishra	96-100

SOCIETY OF BIOLOGICAL SCIENCES AND RURAL DEVELOPMENT**CONTENTS**

➡	SIGNIFICANCE OF ENVIRONMENTAL FACTORS IN HUMAN LIFE Rakesh Kumar Srivastava and Santosh Kumar Arasiya	101-106
➡	CONSTRAINTS AND CHALLENGES IN THE MARKETING Vipul Kumar, N. K. Mishra and Pradeep Kumar Yadav	107-109
➡	A PRELIMINARY STUDY OF PREDATORY BUGS OF MIZORAM, INDIA Sonam Jahan, Preety Khare, Sandeep Kushwaha and Hemlata Pant	110-113
➡	ENTREPRENEURIAL BEHAVIOR OF RURAL WOMEN IN JAUNPUR Ilma Siddique, N.K. Mishra and Pradeep Kumar Yadav	114-118
➡	RURAL DEVELOPMENT THROUGH ENTREPRENEURSHIP Shashi Singh and Kamlesh Singh	119-127
➡	A STUDY ON ADOPTION BEHAVIOUR OF FARMERS ABOUT Mamata Jayasawal and N. K. Mishra	128-132
➡	DEVELOPING AN EFFICIENT AND SUSTAINABLE DISPOSAL Jyoti Verma, Hemlata Pant, Nikita Rawat and Anuradha Yadav	133-141
➡	EFFECT OF LEAD CHLORIDE EXPOSURE ON BLOOD GLUCOSE Amardeep Sagar, Kavita, D. K. Chauhan and Divya Singh	142-145
➡	EFFECT OF NAA AND BORON LEVELS ON PLANT AND CURD Manoj Kumar Singh	146-148
➡	ESSENTIAL OIL: ITS EXTRACTION TECHNIQUES Sujata Gupta, Barkha Kamal, Smita Sharma, Reena Verma and M.K. Purohit	149-155
➡	FISH FAUNA OF JABALPUR DISTRICT OF MADHYA PRADESH Sanjay D. Paunikar, Sandeep Kushwaha and Sangeeta Mashi	156-162
➡	IMPACT OF MASS MEDIA ON RURAL DEVELOPMENT Richa Rai and N. K. Mishra	163-167

A COMPARISON OF OCCURRENCE OF FOSSILS OF TWO WHALE FAMILIES (CETACEA BRISSON, 1762: ODONTOCETI FLOWER, 1867) : KOGIIDAE AND KENTRIODONTIDAE

P.S. Bhatnagar¹ and S. Pujari²

¹Zoological Survey of India

Central Zone Regional Centre, Near SBI Chowk, Jabalpur, Madhya Pradesh - 482002, India

²Govt. College of Arts, Science and Commerce Sanquelim, Goa - 403505, India

(Corresponding author Email: zsiscientist2004@yahoo.com)

Received : 12.11.2022

ABSTRACT

Accepted : 15.12.2022

Kogiidae are a family of small odontocetes ; toothed whales (Cetacea), While on the other hand Kentriodontidae is an extinct family of dolphins and are small to medium sized toothed cetaceans. One reason that Odontocetii are important is that they are highly encephalised; encephalisation next only to humans. As the occurrence of the fossils of two families have not been studied in a comparative way. The present study, therefore, had the objectives of –extracting of data on occurrence of Kogiidae and Kentriodontidae from database www.paleodb.org and comparing the occurrence of different taxa. In family Kogiidae, *Thalassociatus* species appeared first among the Kogiids and its last fossil also occurred the earliest among the Kogiids. While in In Kentriodontidae, *Kentriodon* premix was the oldest in terms of first occurrence. Well preserved records of Kogiidae are very rare , while Kentriodontidae has relatively common fossils. Both the extant species of Kogiidae are in sub-family Kogiinae and are covered in CITES.

Keywords : *Kogiidae, kentriodontidae, fossils, occurrence*

INTRODUCTION

Kogiidae are a family of small odontocetes ; toothed whales (Cetacea), which are similar and closely related to the present day sperm whale (Physeteridae) (Bianucci and Landini,1999). It is rare to get well preserved fossils of Kogiidae. Some of the rare and almost complete skulls described are that by Barnes (1973,1984) and Muizon (1988), the other being by Bianucci and Landini (1999) . *Kogia breviceps* and *K. simus* are the only two living species of this family (Bianucci and Landini,1999).

While on the other hand Kentriodontidae is an extinct family of dolphins and are small to medium sized toothed cetaceans which possibly also had ancestors of the some living species. *Kentriodon hobetsu* from middle Miocene from Hokkaido , Japan was described as a new species on the basis of partial skull , which significantly extended the geographic range of this genus (Ichishima *et al.*,1994). Kentriodontidae has also been described from Miocene sediments from Italy (Bianucci,1994) . They are relatively common fossils and are quite

useful for intercontinental correlations of marine deposits. Dawson (1996) described the family as almost cosmopolitan in distribution.Kentriodontids are small odontocetes with a well-known fossil record in the Miocene. The Kentriodontidae diversified in the Early Miocene and reached peak diversity by the Middle Miocene . Cladistic analysis of Kenriodontidae has also been carried out (Aguirre Fernandez, 2003).

It is obvious from the foregoing review of literature that the occurrence of the fossils of two families have not been studied in a comparative way. The present study, therefore, had the objectives of–

1. Extracting of data on occurrence of Kogiidae and Kentriodontidae from database www.paleodb.org

2. Comparing the occurrence of different taxa.

MATERIALS AND METHODS

The paleo database from www.paleodb.org was used. First, in the analyze section

- i) Count taxa was used
- ii) Thereafter, generate data summary tables was used; in this option a) items to count was chosen as occurrences, and b) fields to tabulate (rows) was selected as 'continent'. The second field (optional) for columns was left blank.
- iii) Thirdly, analysis of taxonomic ranges was used. Taxon name was given and then break taxa into species option was selected. It generated confidence interval taxon list. It was submitted to display confidence interval options, wherein options shown by default were used, as a result of which confidence interval output was obtained. Taxa were ordered by first occurrence.
- iv) The following taxa were excluded from the chart because they could not be mapped to the time scale specified : *Thalassocetus antwerpiensis* (in case of kogiidae).

Table - 1 : Occurence data for family kongiidae fossils

Taxa/ occurrence	First occurrence (Ma)	Last occurrence (Ma)	Confidence interval (Ma)	Number of horizons	Transposition test
<i>Thalassocetus sp.</i>	13.65	11.608	0	1	1
<i>Nanokogia isthmia</i>	11.608	7.246	0	2	0
<i>Scaphokogia cochlearis</i>	11.608	7.246	0	1	1
<i>Kogia sp.</i>	11.608	0.012	26.671	4	0
<i>Praekogia cedrosensis</i>	7.246	5.332	0	2	0
<i>Scaphokogia sp.</i>	7.246	5.332	0	1	1
<i>Aprixokogia kelloggi</i>	5.332	3.6	0	1	1
<i>Kogia prisca</i>	5.332	3.6	0	1	1
<i>Kogia breviceps</i>	5.332	0	1.285	18	0
<i>Kogia pusilla</i>	3.6	2.588	0	1	1
<i>Kogia sima</i>	3.6	0	5.274	5	0

Table - 2 : Occurrence data for family ketrionodontiadae fossils

Taxa/ occurrence	First occurrence (ma)	Last occurrence (ma)	Confidence interval (ma)	Number of horizons	Transposition test
<i>Incacetus broggii</i>	23.03	20.43	0	1	1
<i>Kentriodon pernix</i>	23.03	13.65	9.99	6	0
<i>Kentriodon sp.</i>	23.03	11.608	3.164	16	0
<i>Kampholophos sp.</i>	20.43	15.97	0	1	1
<i>Rudicetus</i>	20.43	15.97	0	2	0
<i>squalodontoides</i>					
<i>Delphinodon sp.</i>	20.43	13.65	32.768	3	0
<i>Delphinodon dividum</i>	20.43	11.608	12.924	5	0
<i>Delphinodon wymanii</i>	15.97	13.65	0	2	0
<i>Hadrodelphis calvertense</i>	15.97	13.65	11.213	3	0
<i>Hadrodelphis poseidon</i>	15.97	13.65	0	2	0
<i>Kentriodon diusinus</i>	15.97	13.65	0	1	1
<i>Kentriodon obscurus</i>	15.97	13.65	0	1	1
<i>Delphinodon carniolicus</i>	15.97	11.608	0	2	0
<i>Kentriodon fuchsii</i>	15.97	11.608	2.521	9	0
<i>Liolithax kernensis</i>	15.97	11.608	10.033	4	0
<i>Lophocetus pappus</i>	15.97	11.608	21.082	3	0
<i>Hadrodelphis sp.</i>	15.97	3.6	59.784	3	0
<i>Belonodelphis peruanus</i>	13.65	11.608	0	1	1
<i>Heterodelphis croatica</i>	13.65	11.608	0	1	1
<i>Heterodelphis leiodontus</i>	13.65	11.608	0	1	1
<i>Kampholophos serrulus</i>	13.65	11.608	0	1	1
<i>Kentriodon hoepfneri</i>	13.65	11.608	0	1	1
<i>Kentriodon schneideri</i>	13.65	11.608	0	1	1
<i>Leptodelphis</i>	13.65	11.608	0	1	1
<i>stavropolitanus</i>					
<i>Macrokentriodon morani</i>	13.65	11.608	0	1	1
<i>Microphocaena podolica</i>	13.65	11.608	0	1	1
<i>Rudicetus sp.</i>	13.65	11.608	0	1	1
<i>Sarmatodelphis</i>	13.65	11.608	0	1	1
<i>moldavicus</i>					
<i>Sophianacetus</i>	13.65	11.608	0	1	1
<i>commenticius</i>					
<i>Tagicetus joneti</i>	13.65	11.608	0	1	1
<i>Liolithax sp.</i>	13.65	7.246	0	2	0
<i>Heterodelphis klinderi</i>	11.608	7.246	0	1	1
<i>Lophocetus calvertensis</i>	11.608	7.246	0	1	1
<i>Lophocetus repenningi</i>	11.608	7.246	0	1	1
<i>Macrokentriodon sp.</i>	11.608	7.246	0	2	0
<i>Pithanodelphis sp.</i>	7.246	5.332	0	1	1

RESULTS AND DISCUSSION

The outcome of the above methods has been depicted in Table 1 (Kogiidae) and Table 2 (Kentriodontidae). In family Kogiidae, *Thalassocetus* species appeared first among the

Kogiids and its last fossil also occurred the earliest among the Kogiids (Table 1). In Kentriodontidae, *Kentriodon premix* was the oldest in terms of first occurrence followed by *Rudicetus squalodontoides*, with *Kentriodon diusinus* and *Kentriodon*

obscurus occurring at the same time after it and further followed by *Kampholophos serrulus* and *Kentriodon schneideri* at the same time (Table 2). This is based on phylogenetic tree by Predo *et al.* (2018).

Most of the taxa had very good Confidence Interval except *Kogiia* species, where further investigation of fossil specimen may be insightful. The taxa occurrence of Kogiidae in Table 1 is broadly in rhyme with phylogeny tree of (Velez-Juarbe, 2015). However, *Kogia priscalis* listed in Table 1 has been considered as *nomen dubium* (Bianucci and Landini, 1999). As per Muizon (1988, 1991), Kogiidae are divided into two subfamilies on the basis of differential architecture of supracranial basin: the Kogiinae (*Kogia* and *Praekogia*) and Scaphokogiinae (*Scaphokogia*). Both the extant species of Kogiidae are in sub-family Kogiinae and are covered in CITES (www.cites.int). Vale-Juarbe (2015) has drawn phylogenetic trees that include Kogiidae. In this phylogenetic tree also, both the extant species of Kogiidae form a single clade. As per these authors, divergence between extant species of *Kogia* occurred at around the late Pliocene, later than previously predicted by molecular estimates. Phylogenetically, an old separation (at least in the Lower Miocene) of Kogiidae and Physteridae is suggested (Bianucci and Landini, 1999).

Ichishima *et al.* (1994) has given a classification of kentriodontidae, which is mainly based on Barnes (1985a). They indicated that Kentriodontines are plausible ancestors to some Delphinidae and mentioned the need of cladistics analysis for Kentriodontidae. All of modern Delphinoid families evolved from middle or late Miocene Kentriodontidae (Barnes, 1985 b). It is widely accepted that living delphinoids (Delphinidae, Phocoenidae and Monodontidae) originated from within Kentriodontidae (Aguirre

Fernandez, 2003)

Predo *et al.* (2018) while describing a new Kentriodontid from early Miocene of Washington state, *Wimahl chinookensis* (one of the oldest known Kentriodontids) has given a phylogenetic tree of kentriodontidae and their review of this family includes only 4 genera; *Wimahl*, *Kampholophos*, *Rudicetus* and *Kentriodon* in this family. They refer to 'Kentriodontids' outside of this clade as stem Delphinidians. As *Wimahl chinookensis* is one of the oldest known Kentriodontid (Predo *et al.*, 2018), so as per Table 2, *Kentriodon permix* would be second most oldest in terms of first and last occurrence.

It may be mentioned that Odontoceti (which includes Kogiidae and Kentriodontidae) are an important group, one of the reasons being that they are highly encephalised; the encephalisation being next only to modern human and larger than all other mammals. Besides, they have behavioural faculties that have been ascribed to humans and to some extent greater apes. (Marino *et al.*, 2004). So, they are important in understanding evolution of brain. However, well-preserved records of Kogiidae are very rare: only two almost complete skulls from late Miocene sediments of Mexico (Barnes, 1973, 1984) and Peru (Muizon, 1988) have been described previously. The third one by Bianucci and Landini (1999). The lack of substantiated kogiid records until the Upper Miocene is probably due to the rarity of these cetaceans (Bianucci and Landini, 1999). On the other hand, fossils of kentriodontidae are relatively common fossils and quite useful for intercontinental correlations of marine deposits (Ichishima *et al.*, 1994).

ACKNOWLEDGEMENTS

Access to the database www.paleodb.org is gratefully acknowledged. The author is grateful to The Director, Zoological Survey of India.

REFERENCES

1. Aguirre Fernández, G. (2013). Neogene dolphins from New Zealand: implications for the evolution of the family Kentriodontidae (Thesis, Doctor of Philosophy). University of Otago. Retrieved from <http://hdl.handle.net/10523/4050>
2. Barnes L.G. (1973) - *Praekogia cedrosensis*, a new genus and species of fossil pygmy sperrn whales from Isla Cedros, Baja California, Mexico. Contr. Sci., nat. Hist. Mus. Los Angeles C., 247 : 1-20, Los Angeles.
3. Barnes L.G. (1976) - Outline of eastern North Pacific fossil cetacean assemblages. Syst. Zool., v. 25 (a), pp.321-343, New Haven.
4. Barnes L.G. (1984) - Fossil odontocetes (Mammalia: Cetacea) from the Almejas Formation, Isla Cedros, Mexico. Paleobios, v. 42, pp. 1-46, Berkeley
5. Barnes, L. G. 1985 a. The Late Miocene dolphin *Pithanodelphis* Abel, 1905 (Cetacea: Kentriodontidae) from California. *Contributions in Science, Natural History Museum of Los Angeles County* **367**, 1-27.
6. Barnes, L. G. 1985 b. Evolution, taxonomy and antitropical distributions of the porpoises (Phocoenidae, Mammalia). *Marine Mammal Science* **1**, 149-65.
7. Bianucci, G. and Landini, W. 1999. *Kogia pusilla* from the middle Pliocene of Tuscany (Italy) and a phylogenetic analysis of the family Kogiidae (Odontoceti, Cetacea). *Rivista Italiana di Paleontologia E Stratigrafia* . 105(3) : 445-453.
8. Dawson, S. 1996. *A description of the skull and postcrania of Hadrodelphis calvertense* Kellogg 1966 and its position within the Kentriodontidae (Cetacea: Delphinoidea). *Journal of Vertebrate Paleontology*. **16**(1) : 125-134.
9. Ichishima, H., Barnes, L. G., Fordyce, R. E., Kimura, M. and Bohaska, D. J. (1994), A review of kentriodontine dolphins (Cetacea; Delphinoidea; Kentriodontidae): Systematics and biogeography. *Island Arc*, 3: 486–492. doi: 10.1111/j.1440-1738.1994.tb00127.x
10. Ichishima, H., Barnes, L.G., Fordyce, R.E., Kimura, M. and Bohaska, D.J. 1994. A review of kentriodontine dolphins (Cetacea ; Delphinoidea; Kentriodontidae) ; Systematics and biogeography . *The Island Arc* 3 : 486-492.
11. Muizon C. de .1988 . Les Vertébrés de la Formation Pisco (Pérou). Troisième partie: Les Odontocètes (Cetacea, Mammalia) du Miocène. *Inst. Fr. Etud. Andine*, mem. 78. pp. 1-244. Paris.
12. Muizon C. de .1991. A new Ziphuidae (Cetacea) from the Early Miocene of Washington State (USA) and phylogenetic analysis of the major groups of odontocetes. *Bull. Mus. natn. Hist. nat.*, 12 (3-4): 279-326, Paris
13. Peredo, C. M., M. D. Uhen, and M. D. Nelson. 2018. A new kentriodontid (Cetacea: Odontoceti) from the early Miocene Astoria Formation and a revision of the stem delphinidan family Kentriodontidae. *Journal of Vertebrate Paleontology*. DOI: 10.1080/02724634.2017.1411357.

INTERCROPPING OF SUGARCANE AND CAULIFLOWER TO GET HIGHER PROFIT RATE IN GONDA DISTRICT OF U.P., INDIA

Pradeep Kumar Misra, Manoj Kumar Singh and R L Yadav

Krishi Vigyan Kendra, Mankapur, Gonda - II, (U.P.), India

Acharya Narendra Dev University of Agriculture and Technology, Kumarganj, Ayodhya, (U.P.), India

Received : 21.12.2022

ABSTRACT

Accepted : 05.02.2023

Sugarcane is the main crop of the Majha (Diara) land of Gonda district, Uttar Pradesh covering with 60-65 percent of total area. Most of the farmers were growing sugarcane as sole crop and got average yield 665 to 810q/ha with the Rs 1.81 lakhs/ha/year net return, which is not good for farmers of gonad district. Keeping in viewcauliflower intercropping with autumn sugarcane was promoted through transfer of technology by frontline demonstration. Yadva and Verma (1984) support for transfer of technology. In this system G-75 hybrid cauliflower nursery was sown in First week of September in low tunnel polyhouse in pots with coco pits and after twenty five days field prepared and sugarcane crop was sown through improved trench planting method at the row to row distance 90 cm in trench and cauliflower nursery transplanted on raised bed between two rows of sugarcane at the distance of 45x30 cm. The recommended dose of fertilizer was uses as basal doses (Half dose of Urea and Full doses of P_2O_5 & K_2O) on the bases soil testing. Remaining Urea was used as top dressing in two spilt doses. Cultural practices like weeding irrigation and plant protection activities were done as per needs time to time. Cauliflower crop was harvested in the month march first week and sold in market. At time of harvesting of cauliflower crop all cultural practices like- digging, weeding and irrigation completed in sugarcane crop. In this demonstration cauliflower yield got 225qt/ha and Sugarcane average yield 820 qt/ha. The net return was Rs 364250 /ha and in sole crop was Rs. 171500 which clearly showed that intercropping of Cauliflower with sugarcane increase theincome of farmers in comparison to sole crop of sugarcane.

Keywords : *Intercropping, sugarcane, cauliflower, yield*

INTRODUCTION

Sugarcane (*Saccharum officinarum*) family *Gramineae* (Poaceae) is widely grown crop in India. It provides employment to over a million people directly or indirectly besides contributing significantly to the national exchequer. Sugarcane growing countries of the world lay between the latitude 36.7° north and 31.0° south of the equator extends from tropical to subtropical zones. Sugar cane originated in New Guinea where it has been

known for thousands of years. Sugar cane plants spread along human migration routes to Asia and the Indian subcontinent. Here it cross-bred with some wild sugar cane relatives to produce the commercial sugar cane we know today.

Cultivation of sugarcane in India dates back to the Vedic period. The earliest mention of sugarcane cultivation is found in Indian writings of the period 1400 to 1000 B.C. It is now widely accepted that India is the original home of *Saccharum* species.

Saccharum barberi and Polynesian group of islands especially New Guinea is the centre of origin of *S. officinarum*. It belongs to family Gramineae (Poaceae), class monocotyledons and order glumaceae sub family panicoidae, tribe Andriopogoneae and sub tribe saccharineae. The cultivated canes belong to two main groups: (a) thin, hardy north Indian types *S. barberi* and *S. Sinense* and (b) thick, juicy noble canes *Saccharum officinarum*. Highly prized cane is *S. officinarum*.

Intercropping is the growing of two or more crops simultaneously on the same piece of land such that the period of overlap is long enough to include the vegetative stage (Gomez and Gomez, 1983). Intercropping double cropping and other mixed cropping practices that allow more efficient uses of on farm resources are among the agricultural practices associated with sustainable crop production (NRC, 1983; Tolera, 2003). Intercropping provides year-round ground cover, or at least for a longer period than monocultures, in order to protect the soil from desiccation and erosion. By growing more than one crop at a time in the same field, farmers maximize water use efficiency, maintain soil fertility, and minimize soil erosion, which are the serious drawbacks of monocropping (Hoshikawa, 1991).

Sugarcane is the main commercial crop grown by most farmers of the Majha (diara) land of Gonda district of eastern U.P. About 65-70% is covered by this crop. Farmers are growing sugarcane in autumn, spring and late spring seasons just after wheat harvesting. In autumn season, sugarcane is sown in the months of Sept-Oct. Most of the farmers are choosing autumn sugarcane as their sole crop and they are getting 700-820 qt/ha by using improved trench method. But, due to increasing Labour and inputs cost, cost of cultivation has increased resulting in farmers not getting good returns. The size of land holding is also decreasing day by day in India, which is caused by division and subdivision of land holding. The Government of India wants doubling the farmers income until 2022. For that

purpose the maximum use of land by intercropping of Vegetable production has a potential of enhancing farmer's income. Keeping this issue in view, the Krishi Vigyan Kendra was conducted demonstration on Cauliflower intercropping with sugarcane in autumn season. (Bakhsh et al., 2004) to increase the farmer's income.

The Uttar Pradesh, Bihar, Punjab, Haryana, Uttarakhand, Madhya Pradesh, Rajasthan, West Bengal, Jharkhand, Assam and other north and north eastern state are come in subtropical belt. Among states Uttar Pradesh is the main cane growing state in the country allocating about 2.2 million ha area (43.7%) for cane cultivation and Maharashtra second cane cultivation state with 19.6% cane area.

Small sugarcane growers need not wait until the harvest of sole crop to obtain financial returns. Intercropping of important short duration crop (Cauliflower) with sugarcane through utilization of the present limited land resources would help to sustain sugarcane cultivation and provide interim return to small and marginal farmers.

In the Eastern Uttar Pradesh sugarcane is normally planted in autumn September-October, this planting of sugarcane in variably yields 20-25%. Cauliflower and cane grow as intercrop because there is higher demand of vegetables in the market for whole year. Since the consumption of vegetable is increasing at faster rate in modern time as these are supplementary item in human diet. Sub-tropical region contributes more than 55% area of the sugarcane, however, cane yield and sugar recovery (%) are lower in comparison with tropical India.

MATERIALS AND METHODS

To keeping in view of the increasing farmer's income the Krishi Vigyan Kendra, Mankapur, Gonda II had conducted a demonstration on cauliflower intercrop with sugarcane under Frontline demonstration in the year 2019-2020 and 2020 and 2021 two constructive years. Total 10 farmers were selected in two villages of covering two blocks i.e., Mankapur and Babahnjot. The

variety CO-238 and G-75 were selected for sowing of sugarcane and cauliflower. Cauliflower (Variety G-75 hybrid) seed treated with 0.3 percentage solution and nursery was raising in germination trayfill with cocopit in the first week of September under low tunnel poly house. Sugarcane was planted through improved trench method in furrows at 90 cm apart keeping three budded sets as 30 cm row length in the third week of September. Ahmed et al., (1997) already promoted double row intercropping.

The use of fertilizer:

Sugarcane fertilized N2 180kg/ha, ½ N2 at planting as based dressing + 1/2 in two equal split doses as top dressing after harvesting of cauliflower at proper moisture while intercrop are given one third of total N+ full doses of P2O5 and K2O at their transplanting on second week of October just after sugarcane planting and remaining half N is top dressed after 30 day of transplanting. Cauliflower Nursery transplanted at the distance of 45x30 cm between two rows of sugarcane on raised bed. Alam et al., (2000) supported these cultural practices done as per need time to time. Observations are given below.

RESULTS AND DISCUSSION

In table No. 2 data shows that sowing time and method of sugarcane planted were same. Seed

requirement of sugarcane with cauliflower inter cropping is 75q/ha and it is also 75q/ha in sole crop but cost of cultivation with intercropping became Rs. 1.65 lakh/ha and Rs 0.98 lakh/ha is in sole crop. Its increase 68.36% in comparison to sole crop. Production of sugarcane is 870qt/ha in intercropping cauliflower yield 232 qt/ha. Sugarcane yield in sole crop was 930qt/ha. Geetha et al. (2015) also worked on yield performance of sugarcane in intercropping.

Total return in Cauliflower intercropping is Rs 5.51 Lakhs/ha and Rs 2.79 Lakhs/ha in sole crop. The total return is 1.97% is excess in comparison to sole crop of sugarcane. The net return is more (Rs 2.72 Lakhs) in comparison to sole crop of sugarcane (Rs 2.79 Lakhs) Ahmed et al., (1997) supported the intercropping system. The benefit cost rate on 2.33 is significant higher in comparison to sole crop (1.84) (Ali and Abedullah, 2002). Data of net return revealed that cauliflower intercropping with spring sugarcane play the significant role toward doubling the income of the farmers. Residual of cauliflower also increased the organic matter in the soil and increased the sustainability of the soil.Priscilla and Singh (2015) also found the same trend in vegetable intercropping with sugarcane in Manipur, India.

Table - 1 : Details of Cauliflower intercropping with autumn sugarcane.

Name of crop	Variety	Row to row distance (cm)	Sowing duration	Plant to plant distance (cm)	Seed rate/ha	Application of fertilizers (Kg/ha)		
						N	P	K
Sugarcane (Main crop)	CO-0238	90	3 rd week of September	30	75 gm	180	60	50
Cauliflower (Intercropping)	G-75 (Hybrid)	45	2 nd week of October	30	350 gm	80	40	40

The same results were also observed by Lithourgidis et al., (2011) during their study. Cauliflowers intercropping with autumn sugarcane also promote and help to the maximum utilization of inputs, soil and water etc.

CONCLUSION

On the Object side, it is concluded in the present investigation cauliflower inter cropping with autumn sugar cane data proves that Cauliflower intercropping with autumn sugarcane plantation is

Table - 2 : Cauliflower intercropping with autumn sugarcane.

Sl. No.	Details	Cauliflower intercropping with spring sugarcane	Sole crop of sugarcane
1	Main crop	Sugarcane (Co-238)	Sugarcane (Co-238)
2	Intercrop	Cauliflower (GS-75))	-
3	Sowing time	Sugarcane-3 rd week of September Cauliflower-2 nd week of October	Sugarcane-3 rd week of September
4	Seed rate	Sugarcane-75qt/ha Cauliflower-350gm/ha	Sugarcane-75qt/ha
5	Cost of cultivation	Rs.- 1.55 Lakh/ha	Rs.- 0.95 Lakh/ha
6	Mill able		
7	Production	Sugarcane-770 qt/ha Cauliflower –225 qt/ha	Sugarcane-820 qt/ha
8	Total Return	Sugarcane-Rs. 250250 Cauliflower-Rs. 270000 Total= Rs. 520250	Rs- 266500
9	Net return	Rs- 364250	Rs-171500
10	C:B Ratio	1:2.35	1:1.81

Note :- Sugarcane @Rs 325/qt, Cauliflower @Rs 1200/qt

profitable (Rs 3.64 Lakhs) in comparison to sole crop of sugarcane cultivation (Rs-1.71Lakhs),the B:C ratio was also Higher (2.35) in intercropping of cauliflower in comparison to the sole crop (1.81) and increase the sustainability of soil. Thus Cauliflower intercropping with autumn sugarcane is recommended for increasing the income of the farmers (Ali and Abedullah, 2002 and Ahmad et al. 2004)autumn plantation of sugar cane should be promoted for increasing income of farmers. With this system inputs and natural resources like soil and water also proper utilized. The input use efficiency also increased. Singh et al. (2015) also worked on vegetables intercropping with autumn planted sugarcane: a step towards doubling farmers' income in Indian sub-tropics. Tripathi , janhavi and Pant, Hemlata, (2012)

REFERENCES

- Ahmad, B. K., Bakhsh and Hassan, S. 2004. Economics of growing different summer vegetables. Faculty of Agricultural Economics and Rural Sociology, University of Agriculture, Faisalabad, Pakistan.
- Ahmed, M. S., Cheema, M. S. and Muhamonad, G. 1997. Feasibility of intercropping Rabi crops in autumn crop of Sugarcane Pak. Sugar J., 5.
- Alam MJ, Rahman MM, Zaman AKMM.2000. Impact of paired row Sugarcane with double intercrops. Bangladesh Journal of Sugarcane,; 22:1-9.
- Ali, M. and Abedullah. 2002. Nutritional and Economics Benefits of Enhanced Vegetable Production and consumption. J. Crop Prod. 6:145- 176.
- Bakhsh, K. W. Akram, M. A. Raza and I. Hassan. 2004. Determination of factors affecting cauliflower cultivation. Int. J. Agric. Biol. 2: 36- 38.
- Chogatapur SV, Deepa GS, Chandranath HT.2017. Intercropping in Sugarcane (Saccharum officinarum L.): A Review. Int. J Pure App. Biosci. 5(2):319-323.
- Geetha P, Sivaraman K, Tayade AS,

- Dhanapal R. 2015. Sugarcane based intercropping system and its effect on cane yield. *Journal of Sugarcane Research*. 5(2):110.
7. Gomez, A. A. and K. A. Gamez. 1983. Multiple cropping in the humid tropics of Asia Ottawa. 32p.
 8. Hoshikawa, K. 1991. Significance of legumes crops in intercropping the productivity and stability of cropping system. Pp- 173-176. In: C. Johanson, K. K. Lee and K. L. Saharawat, (eds.). Phosphorous nutrition of grain legume in the srid tropics. ICRISAT.
 9. Lithourgidis, A. S. Dordas, C. A. Damalas, C. A. 2011. Annual intercrops: an alternative pathway for sustainable agriculture. *Australian Journal of Crop Science*. 5 (4): 396-410.
 10. NRC (National Research Committee). 1983. Sustainable agriculture and the environment in the humid tropics. National Academy Press. Washington D. C., 702 p.
 11. Priscilla, L. and Singh, S. P. 2015. Economics of vegetable production in Manipur, *Indian Journal of Economics and Development*. 11(4):933-938.
 12. Singh SN, Singh P, Rai RK, Pathak AD. 2018. Vegetables intercropping with autumn planted sugarcane: a step towards doubling farmers' income in Indian sub-tropics. *Indian Farming*. 68(1):65-68.
 13. Tolera, A. 2003. Effect of nitrogen, phosphorus farmyard manure and population of climbing bean on the performance of maize (*zea mays* L.) climbing bean (*phasealus vulgaris* L.) intercropping system in Alfisols of Baka. An M.Sc. thesis presented in the school of Graduate studies od Alemaya University. 1-75 p.)
 14. Yadvu, R. L. and Verma, R. P. 1984. Transfer of the intercropping technique to sugarcane growers. *Indian Sugar Crops J.*, 10:1-2



Intercropping cropping of Cauliflower in Sugarcane at Purechain kunwar village, block Rupaideeh Dist. Gonda

AN ECO-FRIENDLY APPROACH TO BIOREMEDIATION OF PESTICIDE FOR THE ENVIRONMENT

Barkha Kamal¹, Sujata Gupta^{2*}, Reena Verma³, M. M. Juwantha⁴ and M. K. Purohit⁵

¹Department of Botany, D.B.S. P.G. College, Dehradun, (U.K.), India

^{2,4}Department of Zoology, D.A.V. P.G. College, Dehradun, (U.K.), India

³Department of Botany, D.A.V. P.G. College, Dehradun, (U.K.), India

⁵Department of Zoology, S.G.R.R. College, Dehradun, (U.K.), India

Corresponding Author: sujatadavpg@gmail.com

Received : 25.11.2022

ABSTRACT

Accepted : 27.12.2022

Bioremediation of pesticide is eco- friendly approach which is based on the assumption that various organisms will interact to eliminate waste compounds or pollutants (pesticide) from the environment (biodegrade). Bacteria and fungi are excellent degraders of complex compounds, which they need to fuel their metabolism and growth. The degradation of wastes or contaminants is not the sole aspect of bioremediation. The two methods of *in situ* bioremediation and *ex situ* bioremediation i.e. Bioventing, Biosparging and Bioaugmentation may be sufficient to eliminate the waste or more significant treatments. As a result, applying microbial enzyme directly to the contaminant can occasionally get beyond the constraints associated with biological samples. Enzymes for pesticide breakdown could be developed as a bioremediation technique. Another concern with bioremediation is the cost-benefit ratio, or the cost vs the overall environmental impact. When we wish to employ a live organism or product to remove toxins, it is critical. Understanding the genetics and biochemistry of the target microbe can help to overcome some of these restrictions and to obtain the desired effect in a short period of time. In the development of technology for eliminating toxins from actual locations, bioremediation has a better future than other remediation techniques such as incineration, thermal disposition, land farming, and so on.

Keywords : Pesticides, bioremediation, *in situ* bioremediation, *ex situ* bioremediation and pollution.

INTRODUCTION

Any approach utilized to eradicate the negative impacts of contaminants from the environment is referred to as "bioremediation." Pollutants must be eradicated whenever possible, although this is not always practicable; nonetheless, some species may be able to limit or immobilize them. Organisms, for example, can accumulate contaminants and decrease their presence and

impact on the environment, but they do not eradicate them. Such methods, which are now in practice (v.gr. phytoextraction) (Dubchak and Bondar 2019, Tyagi *et. al.*, 2011) should be included in the idea of "bioremediation". Bioremediators are organisms that are capable of bioremediation. For example pesticides like Algicides target Algae, Bactericides targets Bacteria. Fungicide targets fungi etc.

Microbes are used to breakdown pesticides

in situ in bioremediation. An effective bioremediation process necessitates the use of a bacterial strain capable of degrading the biggest pollution to the smallest amount possible. To achieve an acceptable level of pesticide residues or metabolites at a polluted site in a reasonable amount of time, an adequate rate of biodegradation is necessary. The rate of biodegradation in soil is determined by four factors: (i) physiological status of microorganisms, (ii) pesticide or metabolite availability to microorganisms (iii) survival and/or proliferation of pesticide degrading microorganisms at contaminated sites, and (iv) a sustainable population of these microorganisms. Water potential, pH, water potential, Temperature, nutrients, and the amount of pesticide or metabolite in soil could all be limiting factors for pesticide-degrading microorganisms, which have to be investigated further in connection to the overall microbial population and their biochemical activities. Because soil has a unique binding ability for a range of pesticides or metabolites, an understanding of pesticide dynamics in the soil environment is also essential when considering an appropriate bioremediation strategy. Before utilizing any bacterium for pollutant biodegradation, it is also vital to understand the genetics and physiology of pollutant degradation. In the terrestrial ecosystem, pesticide contamination can occur in three places: the saturated zone or ground water and the vadose zone or surface soil. Because surface soils have a significant number of aerobic microorganisms, biodegradation is fast and largely aerobic. However, the quantity of aerobic bacteria diminishes with depth. In the other two zones, namely vadose and ground water, biodegradation is slow. To deal with diverse parts of the terrestrial ecosystem, different bioremediation systems are necessary (Nawaz *et al.*, 2015, Tyagi *et al.*, 2011).

Pollution of the environment, freshwater, and topsoil has evolved from global industrialization. Water quality has worsened as a

result of human activity, such as due to mining and ultimate removal of toxic metal effluents from steel mills, battery companies, and electricity generation, posing major environmental concerns. Effluents like petroleum, polythenes, and trace metals harm the environment. Heavy metals are pollutants that exist in nature in the Earth's crust and are difficult to decompose. They exist as ores in rocks and are recovered as minerals. High-level exposures can release heavy metals into the environment. Once in the environment, they remain toxic for much longer (Masindi *et al.*, 2021). Many of these pollutants are mutagenic to both humans along with their surroundings. Absorbing heavy metals accumulates in the brain, liver, and kidney. Other effects on animals include cancer, nervous system damage, stunted growth, and even death (Briffa *et al.*, 2020). Heavy metals in soils reduce food quality and quantity by inhibiting nutrient absorption, plant growth, and physiological metabolic processes. Metal-contaminated soils are being remedied using chemical, biological, and physical methods. However, physicochemical methods produce a lot of waste and pollution, so they are not valued (Rebello *et al.* 2021). Bioremediation is a cost-effective and practical solution for removing environmental contaminants (Tripathi *et al.* 2021).

Bioremediation technology advancement

In the field, bioremediation works by either transforming or degrading the contaminant, or by both methods. In this study, the process technologist's goal is to create the best conditions for microorganism growth. Land farming, bioreactors, biologically enhanced soil washing, composting, and solid-phase bioremediation are all examples of bioremediation treatments. *In situ* bioremediation necessitates the manipulation of aqueous constituents as well as bioventing (Adams *et al.*, 2015). Microbes that have been genetically modified are used to improve their degradation capability. The following points should be considered when developing technology: i) contaminant heterogeneity. (ii) Contaminant

concentration and its effect on biodegradative microbes, (iii) contaminant persistence and toxicity, (IV) contaminant behavior in soil, and (v) conditions favorable to biodegradative microbes or microbial populations.

***In situ* bioremediation**

These techniques are generally the most desirable options because they are less expensive and cause less disruption because they provide treatment in place, avoiding excavation and transport of contaminants. The depth of the soil that can be effectively treated limits *in situ* treatment. Effective oxygen diffusion for desirable rates of bioremediation in many soils extends only a few centimetres to about 30cm into the soil, though depths of 60cm and greater have been effectively treated in some cases. The following are the most important land treatments: The most common *in situ* treatment is bioventing, which involves supplying air and nutrients through wells to contaminated soil in order to stimulate the indigenous bacteria. Bioventing used in low air flow rates, which delivers only the quantity of oxygen required for biodegradation while reducing volatilization and the release of contaminants into the atmosphere. It is effective for simple hydrocarbons and can be employed where contamination is found far beneath the surface. *In situ* biodegradation includes circulating aqueous solutions through polluted soils to provide oxygen and nutrients to naturally occurring microorganisms that breakdown organic pollutant. It's suitable for both soil and groundwater. In general, situations such as the infiltration of water-containing nutrients, oxygen, or other electron acceptors for groundwater treatment are included in this method (Chaney *et al.*, 1997). Biosparging is the process of injecting compressed air beneath the water table to raise groundwater oxygen levels and speed up the biological breakdown of contaminants by naturally occurring microorganisms. Biosparging increases the interaction between soil and groundwater by increasing mixing in the saturated zone.

Bioaugmentation is a term that refers to the use of the addition of microorganisms, either indigenous or exogenous, to contaminated locations is a common part of bioremediation methodology. The utilization of additional microbial cultures in a land treatment unit is limited by two factors: 1) Nonindigenous cultures rarely compete well enough with indigenous populations to develop and sustain viable population levels, and 2) most soils with long-term exposure to biodegradable garbage include indigenous microbes that effectively decompose if the land treatment unit is properly-managed (Adams *et al.*, 2015)

***Ex situ* bioremediation**

These methods entail excavating or removing contaminated dirt from the ground. Landfarming is a basic approach that involves excavating polluted soil, spreading it over a prepared bed, and tilling it periodically until pollutants are reduced. The goal is to encourage indigenous biodegradative bacteria and make it easier for them to degrade pollutants aerobically. Composting is a process that includes mixing polluted soil with non-hazardous organic amendments like manure or agricultural waste. The presence of these organic materials encourages the growth of a diverse microbial population and the high temperatures associated with composting (Philp and Atlas 2005). Landfarming and composting are combined in biopiles. Engineered cells are built similarly to aerated composting piles. They are a refined version of landfarming that tend to control physical losses of contaminants via leaching and volatilization. Biopiles include an ideal environment for aerobic and anaerobic bacteria to thrive. Bioreactors are a type of bioreactor. *Ex situ* treatment of contaminated soil and water pumped up from a contaminated plume is done in slurry reactors or aqueous reactors. In reactor bioremediation, polluted solid material (soil, sediment, sludge) or water is processed through a designed containment system. A slurry bioreactor is a containment tank and apparatus used to create a three-phase (solid, liquid,

and gas) mixing condition in contaminated soil and biomass (typically indigenous) to boost the bioremediation rate of soil-bound and water-soluble contaminants as water slurry. Because the enclosed environment is more manageable and hence more controllable and predictable, the pace and amount of biodegradation in a bioreactor system is larger than *in situ* or in solid-phase systems. Despite the benefits of reactor systems, they do have significant drawbacks. Before being placed in a bioreactor, the contaminated soil requires pretreatment (e.g., excavation), or the contaminant can be removed from the soil via soil washing or physical extraction (e.g., vacuum extraction).

Application of technology at the actual site

The application of technology at the actual site necessitates of natural bioprocesses at the contaminated site, (ii) detailed and valid data of microbial biodegradation developed in the laboratory, and (iii) monitoring of the onsite biodegradation process. Studies on microbial populations, activities, and soil enzymes will provide a mirror image of the soil's functional status. The majority of bioremediation technologies for the field are intended to remove pollutant after it has been generated or released into the environment Bioaugmentation (the addition of an organism or enzyme to a contaminant), biostimulation (the use of nutrients to stimulate naturally occurring organisms), biofilters (the removal of organic gases by passing air through compost or soil containing microorganisms), bioreactors (the treatment of a contaminant in a large tank containing an organism or enzyme), and bioventing (the treatment of a contaminant in a large tank containing an organism or enzyme) are examples of these technologies (Adams *et al.*, 2015, Tyagi *et al.*, 2011)

Bioremediation's Limitations

On the approach to bioremediation technology, there are various challenges. Pollution cleanup by organisms is not a charitable deed. Rather, it is a survival strategy. The majority of bioremediation species work in environments that

are conducive to their demands. As a result, some form of environmental modification is required to induce organisms to decay or absorb the pollution at a rate that is tolerable. The organism must often be exposed to low amounts of the contaminant over a long period of time. This causes the organism to develop the metabolic pathways necessary for the pollutant to be digested. It is frequently necessary to provide fertilizer or oxygen to the substance holding the contaminant when utilizing bacteria and fungi. When done *in situ*, this can be harmful to other creatures. When simple chemicals and metals are taken up, the organisms are likely to be exposed to dangerous quantities of these contaminants. Overall, organisms do not often do as well on a pollutant diet as they do on other nutrients available more frequently in their surroundings. When undertaking *in situ* remediation, this is a concern. Cost/benefit ratios, i.e. cost versus overall impact on the environment, are another obvious source of worry. Neither the government nor the private sector is willing to spend significant sums of money to clean up pollution. Bioremediation is often very expensive, labour intensive, and can take many months to attain acceptable levels of cleanup. Air bioremediation, in particular, is inefficient when compared to the volume of dirty air produced by industry. Another issue is that both *ex situ* and *in situ* technologies have the potential to cause environmental disruption in addition to pollution damage. The long-term consequences of importing naturally occurring non-native bioremediation species into a region remain unknown. Under laboratory conditions, bioremediation has been shown to be effective (Rajendran *et al.*, 2003; Wasi *et al.*, 2011a, b). Bioremediation's popularity is encouraged by the concept that it is more "green" than other remediation procedures. Despite the high risks, companies and individuals are investing in biotechnology futures. As a result, bioremediation companies have a bright future ahead of them, regardless of their long-term efficacy (Fulekar, 2012).

Bioremediation is technology based on the assumption that various organisms will interact to eliminate waste compounds or pollutants (pesticide) from the environment (biodegrade). Bacteria and fungi are excellent degraders of complex compounds, which they need to fuel their metabolism and growth. The degradation of wastes or contaminants is not the sole aspect of bioremediation. However, the method may be sufficient to eliminate the waste or contaminant without degrading in some cases. Bioremediation will function well with some compounds, but it has a number of drawbacks. It is determined by the type of the organisms, the enzyme in question, its concentration and availability, and eventually the microbes' survival. As a result, applying microbial enzyme directly to the contaminant can occasionally get beyond the constraints associated with biological samples. Enzymes for pesticide breakdown could be developed as a bioremediation technique. Another issue with bioremediation is the cost-benefit ratio, or the cost vs the overall environmental impact. When we wish to employ a live organism or product to remove toxins, it is critical. Understanding the genetics and biochemistry of the target microbe can help to overcome some of these restrictions. To obtain the desired effect in a short amount of time, a Super strain can be developed. Multimember microbial consortia can be utilised in the field to accelerate the mineralization of specific pesticides using either elective enrichment or chemostat enrichment. Bioremediation has a brighter future in the development of technologies for removing toxins from actual sites than other remediation procedures such as incineration, thermal disposition, landfarming, and so on (Dua *et al.*, 2002).

REFERENCES

1. Adams, G.O., Fufeyin, P.T., Okoro, S.E. and Ehinomen, I. (2015) 'Bioremediation, biostimulation and bioaugmentation: a review', International Journal of Environmental Bioremediation & Biodegradation, Vol. 3, pp.28–39, doi: 10.12691/ijebb-3-1-5.
2. Dubchak S., Bondar O. (2019) Bioremediation and Phytoremediation: Best Approach for Rehabilitation of Soils for Future Use. In: Gupta D., Voronina A. (eds) Remediation Measures for Radioactively Contaminated Areas. Springer, Cham. https://doi.org/10.1007/978-3-319-73398-2_9
3. Chaney R.L., Malik M., Li Y.M., Brown S.L., Brewer E.P., Angel J.S. and Baker A.J., Phytoremediation of soil metals, Curr. Opin. Biotech., 8, 279-283 (1997)
4. Dua M., Singh A., Sethunathan N. and Johri A., Biotechnology and bioremediation: successes and limitations, App. Microbiol. Biotech., 59(2-3), 143–152 (2002)
5. Fulekar, M. H. (Ed.). (2012). Bioremediation technology: recent advances.
6. Nawaz, K., Hussain, K., Choudary, N., Majeed, A., Ilyas, U., Ghani, A., Lin, F., Ali, K., Afghan, S., Raza, G. and Lashari, M.I. (2011). Eco–friendly role of biodegradation against agricultural pesticides hazards. *Afri. J. Microbiol. Res.*, 5(3): 177–183.
7. Rajendran, P., J. Muthukrishnan and P. Gunasekaran. 2003. Microbes in heavy metal remediation. Indian J. Experimental Bio., 41: 935-944.
8. Tewari, L. Kumar, J. Saini A. (2012) Bioremediation of pesticides by microorganisms: general aspects and recent advances., In book: Bioremediation of Pollutants., International Publishing House Pvt. Ltd. New Delhi (pp.24-29)
10. Tyagi, M, da Fonseca, M.M.R, de Carvalho C.C.C.R (2011) Bioaugmentation and biostimulation strategies to improve the effectiveness of bioremediation processes
11. Biodegradation, Springer 22, pages 231–241.
12. Philp, J.C., Atlas, R.M. (2005) Bioremediation of contaminated soils and aquifers. In: Atlas RM, Philp JC, editors.

Bioremediation: applied microbial solutions for real-world environmental cleanup. Washington: American Society for Microbiology (ASM) Press; 2005. pp. 139–236.

13. Wasi, S., Tabrez, S. and Ahmad, M. (2011a). Suitability of immobilized *Pseudomonas fluorescens* SM1 strain for remediation of phenols, heavy metals and pesticides from water. *Water, Air, and Soil Pollution*, 220(1–4): 89–99.

14. Wasi, S., Tabrez, S. and Ahmad, M. (2011b). Detoxification potential of *Pseudomonas fluorescens* SM1 strain for re-mediation of major toxicants in Indian water bodies. *Water, Air, and Soil Pollution*, 222(1–4): 39–51

15. Masindi, V.; Osman, M.S.; Tekere, M. (2021). Mechanisms and Approaches for the Removal of Heavy Metals from Acid Mine Drainage and Other Industrial Effluents. In *Water Pollution and Remediation: Heavy Metals*; Springer: Berlin/Heidelberg, Germany, pp. 513–537.

16. Briffa, J.; Sinagra, E.; Blundell, R. (2020). Heavy metal pollution in the environment and their toxicological effects on humans. *Heliyon*, 6, e04691.

17. Rebello, S.; Sivaprasad, M.S.; Anoopkumar, A.N.; Jayakrishnan, L.; Aneesh, E.M. (2021). Cleaner technologies to combat heavy metal toxicity. *J. Environ. Manag.*, 296, 113231.

18. Tripathi, M.; Singh, D.N.; Prasad, N.; Gaur, R. (2021). Advanced Bioremediation Strategies for Mitigation of Chromium and Organics Pollution in Tannery. In *Rhizobiont in Bioremediation of Hazardous Waste*; Kumar, V., Prasad, R., Kumar, M., Eds.; Springer: Singapore, 2021; pp. 195–215.

NUTRITIONAL IMPORTANCE AND PROCESSING ASPECTS OF MIRACLE FOOD (MILLETS)

Niraj Kumar, Neeraj Singh and Vivek Singh

Faculty of Engineering & Technology

MGCG University, Chitrakoot, Satna, (M.P), India

Received : 02.10.2022

ABSTRACT

Accepted : 10.12.2022

Millets are a most important food source in arid region including central and western Asia, the Middle East, Africa, North and South America, central and northwest Australia and semi-arid region it includes north and/or south of the deserts (in Africa, Asia and Australia) or inland and at slightly higher elevations (in North America, South America, the Middle East, Africa and Asia). of the world. Millets are good sources of energy. Millet provide us a good source of protein, fatty acids, minerals, vitamins, dietary fiber and polyphenols. They grow in very adverse conditions of climate like low water and high temperatures, no need of fertilizers. They contain all the macronutrients and micronutrients. That's why millets is called Miracle food . Millet contains high amounts of essential amino acids, especially those containing sulfur (methionine and cysteine). Millet processing methods are used to produce a variety of value-added products, and various production methods such as fermentation, germination, peeling, extruding, boiling, puffing, milling, malting, and crushing are used. Processing of millet which can done by milling removes the bran and germ layers that are rich in fibre and phytochemicals & losses in the form of qualitative & qualitative. The millets contains good source of antioxidants, such as phenolic acids and glycated flavonoids. The nutritional significance of millets demands for determination of nutritional characteristics and functional properties of different types of millet crops as well as development of value added products obtained from millets. Processing techniques of millets are involved in making different value added products & in making different techniques are involved such as fermentation, germination, dehulling, extrusion, cooking, puffing, popping, malting, milling, etc. Having a good nutritative value lead in processing which can help the food industry, researchers, and consumers. By adopting suitable processing technique to optimize the nutrient value, increase the bioavailability of nutrients, and help as a nutrition safety concern .

Keywords : Millets, processing, nutrients, dietary fiber, Phytochemical

INTRODUCTION

MILLETS (Facts)

- Origin: Africa and Asia
- Group of annual grasses arid and semiarid regions
- ✍• Small seeded grains
- Gluten free cereals, Non-allergenic, nutritive cereals and superfoods
- ✍• Millet is a grain crop commonly known for its small seeds and high nutritional value. Millet

Millet is a generic term for a group of small-seeded annual grasses grown as grains on marginal lands, primarily in temperate, subtropical and tropical arid regions. Common types of millet available in India are Ragi (finger millet), Jowar (sorghum), Sama (small millet), Bajra (pearl millet) and Variga (froso millet).

India designates 2018 as the Year of Millet. The United Nations General Assembly passed a resolution declaring 2023 as the International Year of the Flag, as India proposed to the Food and Agriculture Organization (FAO). Millets are highly nutritious ,non glutinous, and non acid forming Foods . Millets have enormous potential to form a core component in agriculture whilst offering nutritional and food security benefits. Millet is a staple food in developing countries. Millet is an excellent source of all essential nutrients such as proteins, carbohydrates, fats, minerals, vitamins and bioactive compounds. Millet is an important human food source, and its production has increased in recent decades to meet the nutritional needs of a growing world population. India, Niger and China are the world's largest producers of millet, accounting for more than 55% of world production. India was the world's largest producer of millet. Millet is a grain mainly used for local consumption. It is rich in dietary fiber, protein, vitamins, potassium, phosphorus, iron, magnesium, and other essential minerals that our body needs. As such, it aids in better digestion and has similar nutritional value to fruits and vegetables. Millet has not only health benefits but also nutritional properties and is used in the treatment of health problems such as diabetes, hyperlipidemia, etc.

Millet is also superior to common grains in terms of nutrient composition and nutritional value for modern lifestyles. They are a rich source of dietary fiber, phytochemicals and micronutrients. India is the world's largest producer of millet. India is the world leader in millet production, accounting for about 15% of global production. In India, Rajasthan is the leading producer of millet. More

than 58% of the world's production consists of millet, but few Indians are aware of its health benefits and nutritional value.

Millet: Jowar (millet), Sama (small millet), Ragi (finger millet), Korra (tail millet) and Variga (millet millet). Bajra itself contains a lot of fat, but ragi contains minimal fat. Wheat and rice provide food security, while millet provides many guarantees such as food, health, nutrition, livelihoods, animal feed, etc., making millet an agricultural security product. Millet is used as a food product and is widely available and consumed in rural areas. In rural areas, part of the millet is used for food such as millet and sorghum, and the rest is used as animal feed. Millet plays an important role in traditional diets in many parts of India. Many states in India use different grades of millet. Compared to rice and wheat, all millets are three times more nutritious than rice and wheat. Rice is high in carbohydrates (28%) and low in protein (3%).

It also contains several essential micronutrients such as B vitamins, manganese and iron.

DIMENSIONS OF MILLETS



Nutritional status of millets

MILLET AND ITS POTENTIAL

Millet belongs to a small, rounded annual grass that grows mainly in arid and semi-arid regions. Millet survives on less fertile soils, is drought tolerant, disease tolerant, has a short growing season, and is grown worldwide throughout the year. The word millet dates back to

Table - 1 : Nutritive value of coarse and fine cereals (Per 100 g).

Crop	Protein(g)	Carbohydrate(g)	Fat(g)	Crude fiber(g)	Mineral(g)	Calcium (mg)	Phosphorus (mg)
Sorghum	10.4	72.6	1.9	1.6	1.6	25	222
Pearl millet	11.6	67.5	5.0	1.2	2.3	42	296
Finger millet	7.3	72.0	1.3	3.6	2.7	344	283
Proso millet	12.5	70.4	1.1	2.2	1.9	14	206
Foxtail millet	12.3	60.9	4.3	8.0	3.3	31	290
Kodo millet	8.3	65.9	1.4	9.0	2.6	27	188
Little millet	8.7	75.7	5.3	8.6	1.7	17	220
Barnyard millet	11.6	74.3	5.8	14.7	4.7	14	121

Source: National Institute of Nutrition (NIN), Hyderabad (P Anbukkani et al., 2018).

the beginning of human civilization, and millet is considered the first domesticated grain. Our country, India, is one of the largest producers of pearl millet in the world. Rajasthan is the largest producer of millet. Millet is rich in protein, calcium, dietary fiber and polyphenols, and has unique health benefits, making it an outstanding property.

Types of Millet

Millet differs from other millet in appearance, vegetation, type of grain, degree of maturity, and morphological characteristics.

The main types of millet:

- African millet (Pennisetum glaucum), most commonly consumed by humans.
- Foxtail (Setaria italic),
- Millet or millet (Panicum miliaceum) and
- Pollen millet (Eleusine coracana).
- Minor millets are
- Barnyard millet (Echinochloa spp.),
- Kodo millet (Paspalum scrobiculatum),
- Little millet (Panicum sumatrense),

- Guineamillet (Brachiaria deflexa),
- Browntop millet (Urochloa ramosa),
- Teff (Eragrostis tef),
- Fonio (Digitaria exilis),
- Sorghum (Sorghum spp.) and
- Job's tears (Coix lacrima-jobi)

Need to Process Millet

- Millets were good for us because of their calorie and energy content & millets were a good source of fibre & phytochemicals. Processing involves partial separation and/or modification of three major constituents of millets-germ, starch-containing endosperm and protective pericarp. Millets have processed by features
- Digestibility
- food safety
- organoleptic properties
- ease of convience
- maximize their nutritional value
- meeting demand for pre-processed & convenience food

Structural features of millets

Grain	Type	Shape	Colour
Sorghum	Caryopsis	Spherical	brown White, yellow, red,
Pearl millet	Caryopsis	Ovoid, hexagonal, globose	Grey, white, yellow, brown, purple
Foxtail	Utricle	Elliptical	millet Yellow, pale yellow
Common millet	Utricle	Globose	Light Purple
Finger millet	Utricle	Globose	Yellow, white, red, brown, violet

Source : FAO 1995

Main Objectives

Optimal processing of millet to meet the specific requirements for nutrition for human health and improvement of food quality due to its fine grain size and short shelf life (against pest infestation and rancidity). Millet with skin, such as millet, godo, millet, millet, and millet, is difficult to process. Hulling efficiency (percentage of millet to total processed grain) and product quality are below acceptable levels. Shelf life and packaging of processed millet.

1. Millet - Nutrients

Millet has high nutritional value and is superior to other grains such as wheat and rice in many components. They are an important source of important nutrients such as niacin, magnesium, phosphorus, manganese, iron and potassium. It is rich in protein, dietary fiber, essential amino acids methionine, lecithin and vitamin E. It is an excellent

source of B vitamins, magnesium, antioxidants, manganese, phosphorus and iron. Millet contains 65% carbohydrates, 9% protein, 3% fat, 2-7% crude fiber, vitamins and minerals. Millet is a good source of essential amino acids except for lysine and threonine, but is relatively high in sulfur-containing amino acids such as methionine and cysteine. It is also rich in essential fatty acids such as linoleic, oleic and palmitic acids.

An excellent source of free and bound forms of monogalactosyl, diacylglycerol, digalactosyl diacylglycerol, phosphatidylethanolamine, phosphatidylserine and phosphatidylcholine. other fatty acids. B. Arachic acid, behenic acid and erucic acid are present in trace amounts. Millet oil can be a good source of linoleic acid and tocopherols.

Millet is a gluten-free, alkaline grain. Vitamins such as niacin, folacin, riboflavin, thiamine and phosphorus are found in millet and play an important role in energy production.

Table - 2 : Nutritive value of coarse and fine cereals (per 100 g)

	Protein (g)	Fiber (g)	Minerals (g)	Iron (mg)	Calcium (mg)
Sorghum	10	4	1.6	2.6	54
Pearl millet	10.6	1.3	2.3	16.9	38
Finger millet	7.3	3.6	2.7	3.9	344
Foxtail millet	12. 3	8	3.3	2.8	31
Proso millet	12.5	2.2	1.9	0.8	14
Kodo millet	8.3	9	2.6	0.5	27
Little millet	7.7	7.6	1.5	9.3	17
Barnyard millet	11.2	10.1	4.4	15.2	11

source :- Source: <http://milletindia.org>

II Millet as a healthy diet

Millet is rich in several beneficial nutrients such as phosphorus, magnesium, copper and manganese. Millet can be used to make bread, beer, cereals and other dishes. Millet is a staple food worldwide. A variety of traditional foods and beverages are made of millet, including roti, bread (fermented or unfermented), cereals, snacks and fast food, baby food, millet liquor, and nutritional millet powder.

III. Millets and Phytochemicals

Millets are also rich sources of phytochemicals and micronutrients. Phytochemicals such as phenolics (bound phenolic acid-ferulic acid, free phenolic acid-protocatechuic acid), lignans, β -glucan, inulin, resistant starch, phytates, sterols , tocopherol, dietary fiber and carotenoids are present in millets. The main polyphenols are phenolic acids and tannins, while flavonoids are present in small quantities; they act as antioxidant and play many

roles in the body immune system.

IV. Millets as Probiotic and Prebiotic

Probiotics are live microorganisms that, when ingested in sufficient quantities, exhibit health benefits beyond basic nutrition needs.

Probiotics are viable microbial food supplements that improve host health. Prebiotics can be used as an alternative or additional support to probiotics. Prebiotics have great potential to alter the gut microbiome, but these modifications occur at the individual strain and species level and cannot be easily predicted in advance. Probiotics aid existing flora or help repopulate the colon when bacteria levels are reduced by antibiotics, chemotherapy or illness. Probiotics are "live microorganisms" that, when administered in adequate amounts, confer health benefits to the host.

Fermented millet products act as a natural probiotic remedy for diarrhea in young children. Prebiotics are indigestible food components that exert beneficial effects on the host by selectively stimulating the growth and activity of one or a limited number of bacteria in the large intestine. Prebiotic activity that helps increase the population of beneficial bacteria that play an important role in improving digestion. Malting produces important beneficial biochemical changes in millet grains.

V. Millet as Nutraceutical

Nutraceuticals are natural bioactive compounds that promote health, prevent disease, or have other medicinal properties. Nutraceuticals provide physiological benefits and help maintain good health. A nutraceutical can be defined as a diet. Supplements that provide bioactive food compounds in concentrated form in a non-food matrix to promote health and prevent disease.

Neutraceuticals are the bioactive compounds found in fortified food, dietary supplements, and herbal products. Millets have nutraceutical properties in the form of antioxidants which prevent deterioration of human health such as lowering blood pressure, risk of heart disease, prevention of cancer and cardiovascular diseases,

diabetes, decreasing tumor cases etc. The concepts of food consumption are changing from previous to present time. Previous emphasis has been on survival, hunger satisfaction, health maintenance and absence of adverse effects on health and current emphasis is on encouraging the use of nutraceutical foods which promise to promote better health and well being thus helping to reduce the risk of chronic diseases such as obesity, diabetes, CVD and cancer.

Health Benefits of Millets

Millets are a powerhouse of nutrients.

Millets have therapeutic benefits such as control of asthma, migraine, blood pressure, diabetic heart disease, atherosclerosis and heart attack.

- Millet have a protective effect against insulin resistance, heart disease, diabetes, ischemic stroke, obesity, breast cancer, childhood asthma and premature death. Because of these benefits millets, millets can be used in functional foods and as neutraceuticals

- Millets have many nutraceutical properties that are helpful to prevent many health problems such as lowering blood pressure, risk of heart disease, prevention of cancer and cardiovascular diseases, decreasing tumour cases etc.

Other health benefits are increasing the time span of gastric emptying, provides roughage to gastro intestine . Millet is an alkaline forming food. Alkaline based diet is often recommended to achieve optimal health, meaning when it combines with digestive enzymes. The soothing alkaline nature of millet helps to maintain a healthy pH balance in the body, crucial to prevent illnesses.

- Millets are anti acidic
- Millets are gluten free
- Millets detoxify body
- Niacin (vitamin B3) in millet can help lower cholesterol
- Prevents breast cancer
- Helps to prevent type 2 diabetes
- Effective in reducing blood pressure
- Helps to protect against heart diseases
- Aids in treating respiratory conditions such as

- asthma
- Helps to optimize kidney, liver and immune system health
- Reduces risk of gastrointestinal conditions like gastric ulcers or colon cancer
- Eliminates problems like constipation, excess gas, bloating and cramping
- Millet acts as a prebiotic feeding microflora in your inner ecosystem

CONCLUSION

Millets are staple food source that is not only providing major nutrients like protein, carbohydrate, fat etc.

As well as providing adequate amounts of vitamins and minerals. In developing countries, various health problems such as malnutrition, obesity, diabetes, cardiovascular disease, skin disease, cancer, and celiac disease are the most prominent. This is mainly due to underutilized crops for food and people's ignorance and ignorance. Millets are readily available and inexpensive.

Millets contain many major and minor nutrients like carbohydrate, good protein, fat, dietary fibre, vitamins and minerals as well as antioxidant and phytochemicals. The importance of this study undertakes to concern and developing specific agenda for these crops which must be recognized as an important food and introduce the millets as a nutritious food, fulfillment of the nutritional need of global population and to find ways to consume the millets nutritionally, effectively and to reduce the problems of malnutrition and other health problems. This study focused to reducing some anti-nutrients which diminish the acceptability, digestibility and bioavailability of nutrients and improve the nutrients of millets for nourishing the health. Household food processing strategies are used for improving the nutritional quality to promote millet utilization for future prospective.

This study emphasized on nutraceutical properties of millets and the application of millets as alternative cereals potentially healthy to elaborate

therapeutic food products like protein and energy rich diet, diet for diabetes, gluten free diet, CVD, etc. This study showed that millet is used as a “food medicine”. Millet is a source of antioxidants such as phenolic acids and glycated flavonoids. Millet products are also characterized as potential prebiotics and may increase the viability of probiotics with potential health benefits.

REFERENCES

1. Abd El-Salam MH, Hippen AR, Salem MM, Assem FM & El-Aassar M. Survival of probiotic *Lactobacillus casei* and *Enterococcus fecium* in Domiati cheese of high conjugated linoleic acid content. Emir. J. Food Agric. 2012; 24(2): 98–104.
2. Adekunle AA. Agricultural Innovation in Sub-Saharan Africa: Experiences from Multiple Stakeholder Approaches. Forum for Agricultural Research in Africa, Ghana. 2012; ISBN 978-998-8373-2-4.
3. Devi PB, Vijayabharathi R, Sathyabama S, Malleshi NG, Priyadarisini VB. Health Benefits of Finger Millet (*Eleusine coracana* L.) Polyphenols and Dietary Fiber: A Review.
4. Shahidi F & Chandrasekara A. Millets grain phenolics and their role in disease risk reduction and health promotion: A review. Journal of Functional Foods. 2013; 5: 570-581.
5. Bhattacharjee R, Khairwal IS, Bramel PJ & Reddy KN. Establishment of a pearl millet [*Pennisetum glaucum* (L.) R. Br.] core collection based on geographical distribution and quantitative traits. Euphytica. 2007; 155: 35-45.
6. Obilana AB & Manyasa E. Millets. In: PS Belton & JRN Taylor (Eds.). pp. 177-217. Pseudo cereals and less common cereals: Grain properties and utilization potential. Springer-Verlag: New York. 2002.
7. Veena B. Nutritional, functional and utilization studies on barnyard millet. M. Sc. Thesis, University of Agricultural Sciences, Dharwad (Karnataka), India, 2003.

8. Lue S, Hsieh F, Huff HE. Extrusion cooking corn meal and sugar beet fiber: effects on expansion properties, starch gelatinization, and dietary fiber content. *Cereal Chem.* 1991; 68:227-229.
9. Laminu HH, Modu S & Numan AI. Production, In Vitro Protein Digestibility, Phytate Content and Acceptability of Weaning Foods Prepared from Pearl Millet (*Pennisetum Typhoideum*) and Cowpea (*Vigna Unguiculata*). *Int. J. Nutr. Metabol.* 2011; 3(9): 109–113.
10. Chakraborty SK, Singh DS, Chakraborty S. Extrusion: a novel technology for manufacture of nutritious snack foods. *J Beverage Food World.* 2009; 42:23-26
11. Chandrasekara A & Shahidi F. Bioaccessibility and Antioxidant Potential of Millet Grain Phenolics as Affected by Simulated in Vitro Digestion and Microbial Fermentation. *J. Funct. Foods.* 2012; 4: 226–237.
12. Chandrasekara A & Shahidi F. Content of Insoluble Bound Phenolics in Millets and their Contribution to Antioxidant Capacity. *J. Agric. Food Chem.* 2010; 58: 6706–6714
13. Upadhyaya HD, Gowda CLL, Reddy VG (2007), Morphological diversity in finger millet germplasm introduced from Southern and Eastern Africa, *The Journal of Semi-Arid Tropical Agricultural Research.* 3: 1-3.
14. Singh KP, Mishra A, Mishra HN (2012). Fuzzy Analysis of Sensory Attributes of Bread prepared from millet based composite flours. *LWT-Food Science Technology.* 48:276-82.
15. ICRISAT. International Crops Research Institute for the Semi-Arid, 2007 annual report. 2007.
16. Amadou I, Amza T, Yong-Hui S & Guo-Wei L. Chemical Analysis and Antioxidant Properties of Foxtail Millet Bran Extracts. *Songklanakarin. J. Sci. Technol.* 2011c; 33(5): 509–515.
17. <http://milletindia.org>
18. <http://test1.icrisat.org/Publications/EBooksOnlinePublications/Annulreport2007.pdf>
19. Moreno MDL, Comino I & Sousa C. Alternative Grains of Potential, Raw Material for Gluten-Free Food Development in the Diet of Celiac and Gluten-Sensitive Patients. *Austin. J. of Nutri. and Food Sci.* 2014; 2(3): 9.
20. Rao BR, Nagasampige MH & Ravikiran M. Evaluation of Nutraceutical Properties of Selected Millets. *J. Pharma. Bioallied Sci.* 2011; 3(2): 277-279.
21. Laminu HH, Modu S & Numan AI. Production, In Vitro Protein Digestibility, Phytate Content and Acceptability of Weaning Foods Prepared from Pearl Millet (*Pennisetum Typhoideum*) and Cowpea (*Vigna Unguiculata*). *Int. J. Nutr. Metabol.* 2011; 3(9): 109–113.
22. Gupta N, Srivastava AK & Pandey VN. Biodiversity and Nutraceutical Quality of Some Indian Millets. *Proceedings of the National Academy of Sciences, India Section B: Biological Sci.* 2012; 82(2): 265-273.

BASELINE DATA ON BIRD DIVERSITY IN PRAYAGRAJ AND ADJOINING AREAS OF UTTAR PRADESH, INDIA: IMPLICATIONS FOR CONSERVATION AND MANAGEMENT

Imran Alam*, Md. Noor Alam² and Sandeep Kushwaha³

¹Zoological Survey of India, Prani Vigyan Bhawan,
M Block, New Alipore, Kolkata 700053, West Bengal, India

²Vinoba Bhawe University, Hazaribag, Jharkhand - 8253012, India
Zoological Survey of India,

³Central Zone Regional Center, Jabalpur Madhya Pradesh, India 482002

*Corresponding author: imranalam27@gmail.com

Received : 20.09.2022

ABSTRACT

Accepted : 30.11.2022

This paper provides a comprehensive overview of the avian fauna in Prayagraj, Uttar Pradesh, India, drawing on both primary and secondary data sources. A total of 93 bird species were recorded, belonging to 74 genera, 43 families, and 18 orders. Maximum species were from the Columbidae family, and the maximum birds were recorded from the Sam Higginbottom Institute of Agriculture, Technology, and Sciences (SHI). Out of 93 recorded species, 87 fall under the Least Concern (LC) category of IUCN Red List, four are Near Threatened (NT), and one is Vulnerable (VU). Lesser Adjutant *Leptoptilos javanicus* is the only species in the Vulnerable category among the recorded species. Additionally, the authors report that 91 out of 93 species are protected under the Indian Wildlife (Protection) Act of 1972. Specifically, seven species have been placed under Schedule I, which provides the highest degree of protection under the law. This paper highlights the importance of conservation efforts for the avian fauna in Prayagraj and the significance of laws such as the Indian Wildlife (Protection) Act in achieving this goal.

Keywords : Avian fauna; prayagraj; wildlife protection; endangered species; indian wildlife protection act

INTRODUCTION

Prayagraj (25°27'N 81°51'E), formerly known as Allahabad, is one of the largest cities in India, situated in the northern state of Uttar Pradesh. It is an important cultural, historical, and administrative center of the region. The city is located at the confluence of the Ganga, Yamuna, and Saraswati rivers, which makes it a significant pilgrimage site for Hindus. The confluence point, known as the Sangam, is considered to be one of the holiest places in India.

Prayagraj has a humid subtropical climate,

with hot and dry summers from April to June, followed by warm and humid summers from July to September, and winters from December to February. The city experiences dense fog during the winter season, which can disrupt transportation and cause respiratory problems for residents.

The unique location and climate of Prayagraj support a diverse range of bird species, especially water-dependent species. Although the city does not have any designated Important Bird Areas (IBAs), as per Islam and Rahmani (2004), the Sangam and Ganga ghat attract a significant number

of migratory birds during the winter season. According to the "Fauna of Uttar Pradesh" report published in 2015, the state supports over 500 bird species, while Prayagraj is home to over 300 (ebird.org/india) bird species out of the 1253 species found in India (Praveen et. al., 2023).

The Ganga-Jamuna Doab, of which Prayagraj is a part, is a region that lies between the Ganga and Yamuna rivers and is responsible for the city's unique flora and fauna. The doab is a fertile region that supports agriculture, and the city is known for its production of crops like wheat, rice, and sugarcane.

In addition to its natural resources, Prayagraj is also home to a number of historical and cultural landmarks. The city has a rich heritage of literature, art, and music and is famous for the Kumbh Mela, a Hindu festival that attracts millions of pilgrims from around the world. The festival is held every 12 years in Prayagraj, and during this time, the city becomes a hub of spiritual and cultural activity.

Study Area:

The current study was carried out in Allahabad and adjoining area during the GSDP training programme. The main focus of the study was to train the GSDP students in identifying birds and preparation of familywise checklist with a parallel aim to document the species diversity of the region. During this study we have covered areas such as Botanical Survey of India campus, Allahabad University, Triveni Sangam, Yamuna Ghat, Allahabad Fort, Sam Higginbottom Institute of Agriculture, Technology and Sciences, Minto Park and Company Garden.

MATERIALS AND METHODS

Throughout our trip, we walked over 50 kilometers while conducting surveys from morning till evening, occasionally even during mid-day. Some areas were located far away, requiring additional travel time to reach them. We used 8 X 42 and 8 X 50 binoculars to observe birds, and we made an effort to photograph all of the species that we

encountered. To ensure maximum area coverage and encounter as many species as possible, we visited a new locality every day.

We used the Focal Animal Sampling method, as described by Altmann (1974), to record bird species. This method involves selecting a particular individual or group of individuals and observing their behavior over a specified period of time. By focusing on a particular bird species, we were able to gather detailed information about their behavior and ecology. Our fieldwork allowed us to observe and document a diverse range of bird species in their natural habitats. Through our efforts, we hope to contribute to a better understanding of the ecology and conservation of these fascinating creatures.

RESULTS AND DISCUSSION

The paper provides an updated and comprehensive overview of the avian fauna in Prayagraj, drawing on both primary and secondary data sources. The authors relied on the findings of recent field trips to Allahabad, as well as specimens housed in the National Zoological Collections. Distribution records from Birds of India (Majumder et. al., 2022), and ebird data were also incorporated to supplement the authors' findings. Moreover, the authors examined previous records of species found in earlier literature, which were not present in the National Zoological Collections or not encountered during the authors' field visits. This multi-faceted approach allowed the authors to present a thorough and well-rounded view of the various species that inhabit Prayagraj, Uttar Pradesh.

Parveen et. al., (2022) reported that India is home to 1351 naturally occurring bird species under 26 orders, 114 families and 493 genera. According to the "Fauna of Uttar Pradesh" report published in 2015, the state supports over 500 bird species, while Prayagraj is home to over 300 (ebird.org/india) bird species out of the 1253 species found in India (Praveen et. al., 2023). During this survey a total of 93 bird species were recorded belonging to 74 genera, 43 families and 18 orders. Maximum species

were from Columbidae family. Maximum birds were recorded from Sam Higginbottom Institute of Agriculture, Technology and Sciences (SHI) (fig. 1). As per IUCN Threat category, out of 93 recorded species, 87 falls under the Least Concern (LC) category of IUCN Red list, 4 are Near Threatened (NT) and one is Vulnerable (VU). The four Near Threatened species are Black-headed Ibis *Threskiornis melanocephalus*, River Lapwing *Vanellus duvaucelii*, Oriental Darter *Anhinga melanogaster* and Alexandrine Parakeet *Psittacula eupatria*. Lesser Adjutant *Leptoptilos javanicus* is the only species in Vulnerable category among the recorded species. While taking into consideration the status of Indian Wildlife Protection act, 91 out of 93 species are protected under the law

The Indian Wild Life (Protection) Act, 1972 is divided into six schedules, each providing varying degrees of protection to different plant and animal species. Schedule I and Part II of Schedule II offer the highest level of protection, with the most severe penalties prescribed for any infringements of these schedules. Species listed in Schedule III and Schedule IV are also protected, but the penalties for violations are less severe than those for Schedule I and Part II of Schedule II. Schedule V lists animals that are classified as vermin and may be hunted freely. However, the hunting of animals listed under Schedule V is subject to regulation by the authorities, and fines may be imposed for any violations of the rules. In conclusion, the Wild Life (Protection) Act of 1972 is a crucial piece of legislation that has helped to conserve and protect India's natural resources. Its six schedules provide a framework for the protection of a wide range of plant and animal species, and its enforcement ensures that the law is upheld and any violations are punished.

Regarding the Prayagraj, it is noteworthy that a total of 91 species have been reported from this region. Among these species, it has been reported that they are classified under three different schedules of the Indian Wildlife (Protection) Act,

1972. Specifically, 7 species have been placed under Schedule I, which provides the highest degree of protection under the law. These species are considered to be highly endangered and threatened, and any violations of this schedule result in the most severe penalties. Indian Peafowl *Pavo cristatus*, Lesser Adjutant *Leptoptilos javanicus*, Crested Serpent Eagle *Spilornis cheela*, Western Marsh Harrier *Circus aeruginosus*, Shikra *Accipiter badius*, Small Minivet *Pericrocotus cinnamomeus* and Rufous-fronted Prinia *Prinia buchanani* are Schedule I species that receives highest degree of protection. Aprt from this, 84 are listed under Schedule II of the act which provide relatively less protection than Schedule I. However, these species are still considered to be important and in need of conservation efforts. Any violations of this schedule are punishable by less severe penalties than those for Schedule I.

The findings of this study have important implications for the conservation and management of bird species in Prayagraj, Uttar Pradesh. With 91 species of birds being reported in the region, it is evident that there is a significant diversity of avian fauna in the area. It is encouraging that the vast majority of species recorded are classified under the Least Concern category of the IUCN Red List, indicating that they are not at immediate risk of extinction. However, the presence of four Near Threatened species and one Vulnerable species highlights the need for conservation efforts to protect these species and their habitats. The identification of Schedule I species in the region underscores the importance of stringent conservation measures to prevent further decline in their populations. It is crucial that the government and conservation organizations take proactive steps to protect these species by enforcing the Wildlife (Protection) Act of 1972, implementing habitat restoration programs, and promoting awareness among local communities. The findings of this study provide a basis for the development of targeted conservation strategies to protect the avian fauna of

Prayagraj and ensure their long-term survival.

Species Diversity




Fig. - 1 : Pie-chart showing area wise bird sightings.




Fig. - 2 : Pie-chart showing number of species protected under different Schedules of Indian Wildlife Protection Act, 1972.

Table - 1 : List of Species observed from Prayagraj, Uttar Pradesh.

Sl No	Order	Family	Species	Authority	IUCN Category	WPA Schedule
1	Anseriformes	Accipitridae	Ruddy Shelduck <i>Tadorna ferruginea</i>	(Pallas, 1764)	Least Concern	Schedule-II
2	Anseriformes	Accipitridae	Indian Spot-billed Duck <i>Anas poecilorhyncha</i>	Forster, JR, 1781	Least Concern	Schedule-II
3	Galliformes	Accipitridae	Indian Peafowl <i>Pavo cristatus</i>	Linnaeus, 1758	Least Concern	Schedule-I
4	Galliformes	Accipitridae	Black Francolin <i>Francolinus francolinus</i>	(Linnaeus, 1766)	Least Concern	Schedule-II
5	Galliformes	Accipitridae	Grey Francolin <i>Ortygornis pondicerianus</i>	(Gmelin, JF, 1789)	Least Concern	Schedule-II
6	Columbiformes	Alaudidae	Rock Pigeon <i>Columba livia</i>	Gmelin, JF, 1789	Least Concern	Not protected
7	Columbiformes	Alcedinidae	Oriental Turtle Dove <i>Streptopelia orientalis</i>	(Latham, 1790)	Least Concern	Schedule-II
8	Columbiformes	Alcedinidae	Eurasian Collared Dove <i>Streptopelia decaocto</i>	(Fridvaldszky, 1838)	Least Concern	Schedule-II
9	Columbiformes	Alcedinidae	Spotted Dove <i>Spilopelia chinensis</i>	(Scopoli, 1786)	Least Concern	Schedule-II
10	Columbiformes	Alcedinidae	Asian Emerald Dove <i>Chalcophaps indica</i>	(Linnaeus, 1758)	Least Concern	Schedule-II
11	Columbiformes	Anatidae	Yellow-footed Green Pigeon <i>Treron phoenicopterus</i>	(Latham, 1790)	Least Concern	Schedule-II
12	Pterocliiformes	Anatidae	Chestnut-bellied Sandgrouse <i>Pterocles exustus</i>	Temminck, 1825	Least Concern	Schedule-II
13	Cuculiformes	Anhingidae	Greater Coucal <i>Centropus sinensis</i>	(Stephens, 1815)	Least Concern	Schedule-II
14	Cuculiformes	Apodidae	Sirkeer Malkoha <i>Taccocua leschenaultii</i>	Lesson, 1830	Least Concern	Schedule-II
15	Cuculiformes	Ardeidae	Pied Cuckoo <i>Clamator jacobinus</i>	(Boddaert, 1783)	Least Concern	Schedule-II
16	Cuculiformes	Ardeidae	Asian Koel <i>Eudynamis scolopaceus</i>	(Linnaeus, 1758)	Least Concern	Schedule-II
17	Caprimulgiformes	Ardeidae	Large-tailed Nightjar <i>Caprimulgus macrurus</i>	Horsfield, 1821	Least Concern	Schedule-II
18	Caprimulgiformes	Ardeidae	Indian Nightjar <i>Caprimulgus asiaticus</i>	Latham, 1790	Least Concern	Schedule-II
19	Caprimulgiformes	Bucerotidae	Asian Palm Swift <i>Cypsiurus balasensis</i>	(Gray, JE, 1829)	Least Concern	Schedule-II
20	Gruiformes	Campephagidae	Eurasian Coot <i>Fulica atra</i>	Linnaeus, 1758	Least Concern	Schedule-II
21	Gruiformes	Campephagidae	Grey-headed Swampphen <i>Porphyrio poliocephalus</i>	(Latham, 1801)	Not Recognised	Schedule-II
22	Gruiformes	Campephagidae	White-breasted Waterhen <i>Amaurornis phoenicurus</i>	(Pennant, 1769)	Least Concern	Schedule-II
23	Gruiformes	Caprimulgidae	Ruddy-breasted Crake <i>Zapornia fusca</i>	(Linnaeus, 1766)	Least Concern	Schedule-II
24	Charadriiformes	Caprimulgidae	River Lapwing <i>Vanellus divaucelii</i>	(Lesson, 1826)	Near Threatened	Schedule-II
25	Charadriiformes	Charadriidae	Red-wattled Lapwing <i>Vanellus indicus</i>	(Boddaert, 1783)	Least Concern	Schedule-II
26	Charadriiformes	Charadriidae	Bronze-winged Jacana <i>Metopidius indicus</i>	(Latham, 1790)	Least Concern	Schedule-II
27	Charadriiformes	Ciconiidae	Common Sandpiper <i>Actitis hypoleucos</i>	(Linnaeus, 1758)	Least Concern	Schedule-II
28	Ciconiiformes	Ciconiidae	Asian Openbill <i>Anastomus oscitans</i>	(Boddaert, 1783)	Least Concern	Schedule-II
29	Ciconiiformes	Cisticolidae	Lesser Adjutant <i>Leptoptilos javanicus</i>	(Horsfield, 1821)	Vulnerable	Schedule-I
30	Suliformes	Cisticolidae	Oriental Darter <i>Anhinga melanogaster</i>	Pennant, 1769	Near Threatened	Schedule-II
31	Suliformes	Cisticolidae	Little Cormorant <i>Microcarbo niger</i>	(Vieillot, 1817)	Least Concern	Schedule-II
32	Pelecaniformes	Columbidae	Little Egret <i>Egretta garzetta</i>	(Linnaeus, 1766)	Least Concern	Schedule-II
33	Pelecaniformes	Columbidae	Cattle Egret <i>Bubulcus ibis</i>	(Linnaeus, 1758)	Least Concern	Schedule-II

Sl No	Order	Family	Species	Authority	IUCN Category	WPA Schedule
34	Pelecaniiformes	Columbidae	Indian Pond Heron <i>Ardeola grayii</i>	(Sykes, 1832)	Least Concern	Schedule-II
35	Pelecaniiformes	Columbidae	Black-crowned Night Heron <i>Nycticorax nycticorax</i>	(Linnaeus, 1758)	Least Concern	Schedule-II
36	Pelecaniiformes	Columbidae	Glossy Ibis <i>Plegadis falcinellus</i>	(Linnaeus, 1766)	Least Concern	Schedule-II
37	Pelecaniiformes	Columbidae	Black-headed Ibis <i>Threskiornis melanocephalus</i>	(Latham, 1790)	Near Threatened	Schedule-II
38	Accipitriformes	Coraciidae	Black-winged Kite <i>Elanus caeruleus</i>	(Desfontaines, 1789)	Least Concern	Schedule-II
39	Accipitriformes	Corvidae	Crested Serpent Eagle <i>Spilornis cheela</i>	(Latham, 1790)	Least Concern	Schedule-I
40	Accipitriformes	Corvidae	Western Marsh Harrier <i>Circus aeruginosus</i>	(Linnaeus, 1758)	Least Concern	Schedule-I
41	Accipitriformes	Corvidae	Shikra <i>Accipiter badius</i>	(Gmelin, JF, 1788)	Least Concern	Schedule-I
42	Accipitriformes	Cuculidae	Black Kite <i>Milvus migrans</i>	(Boddaert, 1783)	Least Concern	Schedule-II
43	Strigiformes	Cuculidae	Spotted Owlet <i>Athene brama</i>	(Temminck, 1821)	Least Concern	Schedule-II
44	Bucerotiformes	Cuculidae	Indian Grey Hornbill <i>Ocyeros birostris</i>	(Scopoli, 1786)	Least Concern	Schedule-II
45	Coraciiformes	Cuculidae	Common Kingfisher <i>Alcedo atthis</i>	(Linnaeus, 1758)	Least Concern	Schedule-II
46	Coraciiformes	Dicruridae	Stork-billed Kingfisher <i>Pelargopsis capensis</i>	(Linnaeus, 1766)	Least Concern	Schedule-II
47	Coraciiformes	Dicruridae	White-throated Kingfisher <i>Halcyon smyrnensis</i>	(Linnaeus, 1758)	Least Concern	Schedule-II
48	Coraciiformes	Estrildidae	Pied Kingfisher <i>Ceryle rudis</i>	(Linnaeus, 1758)	Least Concern	Schedule-II
49	Coraciiformes	Hirundinidae	Green Bee-eater <i>Merops orientalis</i>	Latham, 1801	Least Concern	Schedule-II
50	Coraciiformes	Jacanidae	Blue-checked Bee-eater <i>Merops persicus</i>	Pallas, 1773	Least Concern	Schedule-II
51	Coraciiformes	Leiothrichidae	Blue-tailed Bee-eater <i>Merops philippinus</i>	Linnaeus, 1767	Least Concern	Schedule-II
52	Coraciiformes	Megalaimidae	Chestnut-headed Bee-eater <i>Merops leschenaulti</i>	Vieillot, 1817	Least Concern	Schedule-II
53	Coraciiformes	Meropidae	Indian Roller <i>Coracias benghalensis</i>	(Linnaeus, 1758)	Least Concern	Schedule-II
54	Piciformes	Meropidae	Coppersmith Barbet <i>Psilopogon haemacephalus</i>	(Müller, PLS, 1776)	Least Concern	Schedule-II
55	Piciformes	Meropidae	Eurasian Wryneck <i>Jynx torquilla</i>	Linnaeus, 1758	Least Concern	Schedule-II
56	Piciformes	Meropidae	Brown-capped Pygmy Woodpecker <i>Yungipicus nanus</i>	(Vigors, 1832)	Least Concern	Schedule-II
57	Psittaciformes	Motacillidae	Alexandrine Parakeet <i>Psittacula eupatria</i>	(Linnaeus, 1766)	Near Threatened	Schedule-II
58	Psittaciformes	Motacillidae	Rose-ringed Parakeet <i>Psittacula krameri</i>	(Scopoli, 1769)	Least Concern	Schedule-II
59	Psittaciformes	Motacillidae	Plum-headed Parakeet <i>Psittacula cyanocephala</i>	(Linnaeus, 1766)	Least Concern	Schedule-II
60	Passeriformes	Motacillidae	Indian Pitta <i>Pitta brachyura</i>	(Linnaeus, 1766)	Least Concern	Schedule-II
61	Passeriformes	Muscicapidae	Small Minivet <i>Pericrocotus cinnamomeus</i>	(Linnaeus, 1766)	Least Concern	Schedule-I
62	Passeriformes	Nectariniidae	Long-tailed Minivet <i>Pericrocotus ethologus</i>	Bangs and Phillips, 1914	Least Concern	Schedule-II
63	Passeriformes	Oriolidae	Large Cuckooshrike <i>Coracina macei</i>	(Lesson, 1831)	Least Concern	Schedule-II
64	Passeriformes	Oriolidae	Indian Golden Oriole <i>Oriolus kundoo</i>	Sykes, 1832	Least Concern	Schedule-II
65	Passeriformes	Passeridae	Black-hooded Oriole <i>Oriolus xanthornus</i>	(Linnaeus, 1758)	Least Concern	Schedule-II
66	Passeriformes	Phalacrocoracidae	Black Drongo <i>Dicrurus macrocercus</i>	Vieillot, 1817	Least Concern	Schedule-II
67	Passeriformes	Phasianidae	White-bellied Drongo <i>Dicrurus caeruleascens</i>	(Linnaeus, 1758)	Least Concern	Schedule-II
68	Passeriformes	Phasianidae	Rufous Treepie <i>Dendrocitta vagabunda</i>	(Latham, 1790)	Least Concern	Schedule-II
69	Passeriformes	Phasianidae	House Crow <i>Corvus splendens</i>	Vieillot, 1817	Least Concern	Not protected
70	Passeriformes	Picidae	Large-billed Crow <i>Corvus macrorhynchos</i>	Wagler, 1827	Least Concern	Schedule-II
71	Passeriformes	Picidae	Grey-headed Canary-flycatcher <i>Culicicapa ceylonensis</i>	(Swainson, 1820)	Least Concern	Schedule-II
72	Passeriformes	Pittidae	Ashy-crowned Sparrow Lark <i>Eremopterix griseus</i>	(Scopoli, 1786)	Least Concern	Schedule-II
73	Passeriformes	Ploceidae	Common Tailorbird <i>Orthotomus sutorius</i>	(Pennant, 1769)	Least Concern	Schedule-II
74	Passeriformes	Psittaculidae	Rufous-fronted Prinia <i>Prinia buchanani</i>	Blyth, 1844	Least Concern	Schedule-I
75	Passeriformes	Psittaculidae	Ashy Prinia <i>Prinia socialis</i>	Sykes, 1832	Least Concern	Schedule-II
76	Passeriformes	Psittaculidae	Wire-tailed Swallow <i>Hirundo smithii</i>	Leach, 1818	Least Concern	Schedule-II
77	Passeriformes	Pteroclididae	Red-vented Bulbul <i>Pycnonotus cafer</i>	(Linnaeus, 1766)	Least Concern	Schedule-II
78	Passeriformes	Pycnonotidae	Red-whiskered Bulbul <i>Pycnonotus jocosus</i>	(Linnaeus, 1758)	Least Concern	Schedule-II
79	Passeriformes	Pycnonotidae	Jungle Babbler <i>Argya striata</i>	(Dumont, 1823)	Least Concern	Schedule-II
80	Passeriformes	Rallidae	Brahminy Starling <i>Sturnia pagodarum</i>	(Gmelin, JF, 1789)	Least Concern	Schedule-II
81	Passeriformes	Rallidae	Chestnut-tailed Starling <i>Sturnia malabarica</i>	(Gmelin, JF, 1789)	Least Concern	Schedule-II
82	Passeriformes	Rallidae	Common Myna <i>Acridotheres tristis</i>	(Linnaeus, 1766)	Least Concern	Schedule-II

Sl No	Order	Family	Species	Authority	IUCN Category	WPA Schedule
83	Passeriformes	Rallidae	Bank Myna <i>Acridotheres ginginianus</i>	(Latham, 1790)	Least Concern	Schedule-II
84	Passeriformes	Scolopacidae	Jungle Myna <i>Acridotheres fuscus</i>	(Wagler, 1827)	Least Concern	Schedule-II
85	Passeriformes	Stenostiridae	Oriental Magpie Robin <i>Copsychus saularis</i>	(Linnaeus, 1758)	Least Concern	Schedule-II
86	Passeriformes	Strigidae	Purple Sunbird <i>Cinnyris asiaticus</i>	(Latham, 1790)	Least Concern	Schedule-II
87	Passeriformes	Sturnidae	Baya Weaver <i>Ploceus philippinus</i>	(Linnaeus, 1766)	Least Concern	Schedule-II
88	Passeriformes	Sturnidae	Scaly-breasted Munia <i>Lonchura punctulata</i>	(Linnaeus, 1758)	Least Concern	Schedule-II
89	Passeriformes	Sturnidae	House Sparrow <i>Passer domesticus</i>	(Linnaeus, 1758)	Least Concern	Schedule-II
90	Passeriformes	Sturnidae	Grey Wagtail <i>Motacilla cinerea</i>	Tunstall, 1771	Least Concern	Schedule-II
91	Passeriformes	Sturnidae	Western Yellow Wagtail <i>Motacilla flava</i>	Linnaeus, 1758	Least Concern	Schedule-II
92	Passeriformes	Threskiornithidae	White Wagtail <i>Motacilla alba</i>	Linnaeus, 1758	Least Concern	Schedule-II
93	Passeriformes	Threskiornithidae	Paddyfield Pipit <i>Anthus rufulus</i>	Vieillot, 1818	Least Concern	Schedule-II

ACKNOWLEDGEMENTS

The authors are grateful to the Director, Zoological & Botanical Survey of India, Kolkata for providing necessary facilities and encouragement for preparing the manuscript. We are thankful to Dr. R. P. Sinha, Officer-in-Charge, Central Regional Centre of BSI for logistic and financial support as well as laboratory facilities at Allahabad. We are grateful to Dr. Sandeep Malhotra, Professor of Zoology for his support during GSDP programme and field visit. We are thankful all the GSDP trainees and resource persons for their support during the field work. This work was supported by the Ministry of Environment, Forest and Climate Change (MoEF&CC), Government of India.

REFERENCES

1. Altman. J., 1974, Observational Study of Behaviour: Sampling Methods. Vol. 49, No. 3/4 (1974), pp. 227-267.

2. *Fauna of Uttar Pradesh, State Fauna Series*, 22 (Part - 1): 61-109, 2015. Published by Director Zoological Survey of India.

3. Islam, M. Z. and Rahmani, A. R. (2004) *Important Bird Areas in India: Priority sites for conservation*. Indian Bird Conservation Network: Bombay Natural History Society and BirdLife International (UK). Pp. xviii + 1133.

4. Majumder, A., Maheswaran G., Alam I., Chandra, K., Alfred, J.R.B., Roy Chowdhury, B. 2022. *Birds of India* 1-600. Published by Zoological Survey of India. ISBN: 978-81-8171-581-4

5. Praveen J., & Jayapal, R., 2022. Checklist of the birds of India (v6.2).

6. Praveen J., Jayapal R., & Pittie A., 2016. A Checklist of Birds of India. *Indian BIRDS* 11 (5&6): 113-172.

EXPLORING CONSTRAINTS FACED BY FARMERS IN THE ADOPTION OF MUSHROOM PRODUCTION TECHNOLOGY

Anish Kumar and N. K. Mishra

Department of Agricultural Extension

T. D. P.G. College, Jaunpur, (U.P.), India

Corresponding email : anish8084kr@gmail.com

Received : 03.12.2022

ABSTRACT

Accepted : 11.01.2023

The adoption of mushroom production technology in rural areas is crucial for improving the livelihoods of farmers. However, there are several constraints that hinder the adoption of this technology. This study aims to explore the constraints faced by farmers in the adoption of mushroom production technology in District Nalanda, Bihar. The study found that the major constraints faced by mushroom growers were heavy domestic workload, which was reported by 80.80% of the respondents. This was followed by distance problem, as the Krishi Vigyan Kendra (KVK) was far from the village was reported by 75.8% of the respondents, less freedom of movement (69.2%) and little time available to spend away from home (58.3%).

***Keywords :** Explore, freedom of movement, livelihoods, production technology*

INTRODUCTION

The study was conducted in Nalanda region of Bihar. The KVK of Harnaut Nalanda Bihar was selected for the study. This KVK was operated by Rajendra Agricultural University Pusa, Bihar. In the Impact of training program on knowledge and Adaptation Level for Mushroom, the study was conducted to assess the functioning and effectiveness of selected KVKs in Nalanda Region of Nalanda Bihar. The finding of the study was focused on the KVK in Nalanda Region of Bihar under different system such as Rajendra Agricultural University. It will highlights the problem as well as the necessary steps required to ameliorate funds which had invested on the Mushroom training programme by the central and state government, but very little systematic and planned attempt had been made to evaluate their

effectiveness. For this purpose, training centre had set up all over the country stated earlier, when training programmes of greater magnitude had undertaken, it called for the scientific research with reference to each aspect of the training programme. The evaluation of performance of KVKs regarding training programme and other extension activities was essential to improve the content, quality and methodology. The scheme of KVKs was operating since a long time. The time to assess the performance was made systematically. The study had combined quantitative and qualitative approaches to enable an insight working of KVKs and the reactions of the KVKs functionaries and beneficiaries.

Mushroom production technology has the potential to provide sustainable livelihood opportunities to small farmers in rural areas.

However, the adoption of this technology is often hindered by various constraints. In order to promote the adoption of mushroom production technology, it is important to understand the constraints faced by farmers in its adoption. This study aims to explore the constraints faced by farmers in the adoption of mushroom production technology in District Nalanda, Bihar. District Nalanda is one of the major agricultural districts of Bihar, where mushroom cultivation is gaining popularity as an alternative source of income for small farmers. The Krishi Vigyan Kendra (KVK) and other government agencies have been promoting mushroom production technology in this district through training programs and other extension services. Despite these efforts, the adoption of mushroom production technology is still low among farmers.

MATERIALS AND METHODS

The selected farmers were interviewed using a structured schedule to collect data on the constraints they faced in the adoption of mushroom production technology. The schedule was developed based on a review of relevant literature and expert opinion. The data collected through the schedule was analyzed using descriptive statistics such as frequency distribution and percentage analysis.

RESULTS AND DISCUSSION

Farming Experience:

It is observed from Table 1 that the majority of the respondents i.e. 56.7 per cent were belonged to below 10 years of farming experience followed by 29.2 per cent of the respondents were between 10-25 years of farming experience and 14.2 per cent of the respondents were above 25 years of farming experience.

Table - 1 : Farming experience wise distribution of respondents.

S.N.	Farming Experience	Frequency	Percentage
1.	Below 10 years	68	56.7
2.	Between 10-25 years	35	29.2
3.	Above 25 years	17	14.2
	Total	120	100.00

Thus, it is concluded that the high level of farming experience of respondents belonged to above 10 years of farming experience. The findings of the present study were in accordance with the findings of Thoke and Gunjal (2010).

Farm Holding:

It was observed from Table 2 that the majority of the respondents 53.3 per cent were belonged to small farmer group followed by 36.3 per cent of the respondents were belonged to marginal farmer group whereas, 10.0 per cent were belonged to large farmer group.

Table - 2 : Size of land holding wise distribution of the respondents.

S.N.	Farm Holding	Frequency	Percentage
1.	Small farmer	64	53.3
2.	Marginal farmer	44	36.3
3.	Large farmer	12	10.0
	Total	120	100.00

Thus, it was concluded that the maximum number of respondents were belonged to small farmer group. The findings of the present study were in accordance with the findings of Patil (2008).

Annual income wise distribution of the respondents

It was cleared from Table 3 that, majority 68.3 per cent of the respondents were in the medium annual income category, 16.7 per cent of the respondents were from high annual income category and 15 per cent of the respondents were from low income category.

Table - 3 : Annual income wise distribution of the respondents.

S.N.	Annual Income	Frequency	Percentage
1.	Low (Below Rs 40000)	18	15.0
2.	Medium (Between Rs. 40000-Rs. 80000)	82	68.3
3.	High (Above Rs 80000)	20	16.7
	Total	120	100.00

It was inferred from the present study that the majority of respondents had medium annual income category. The findings of the present study were in accordance with the findings of Anchule (2000).

Social Participation:

The data presented in Table 4 revealed that majority of the respondents of 65.8 per cent belonged to medium social participation category followed by 20.8 per cent of the respondents belonged to low social participation category and 13.3 per cent of the respondents were under high social participation category.

Table - 4 : Distribution of respondents according to their social participation

S.N.	Social Participation	Frequency	Percentage
1.	Low (below 6)	25	20.8
2.	Medium (Between 6-9)	79	65.8
3.	High (Above 9)	16	13.3
	Total	120	100.00

Thus, it was concluded that the maximum respondents were belonging to medium social participation category. The findings of the present study were in accordance with the findings of Biswas *et al.* (2008).

Participation in Extension activities:

The data presented in Table 5 revealed that majority of the respondents of 62.5 per cent belonged to medium category followed by 21.7 per cent of the respondents belonged to high category and 15.8 per cent of the respondents were under low category.

Table - 5 : Participation in extension activities wise distribution of the respondents

S.N.	Participation in Extension	Frequency	Percentage
1.	Low (below 6)	19	15.8
2	Medium (Between 6-9)	75	62.5
3.	High (Above 9)	26	21.7
	Total	120	100.00

Thus, it was concluded that the maximum

respondents were belonging to medium category. The findings of the present study were in accordance with the findings of Biswas *et al.* (2008).

Mass Media Participation:

The data presented in Table 6 revealed that majority of the respondents of 42.5 per cent belonged to medium category followed by 39.2 per cent of the respondents belonged to high category and 18.3 per cent of the respondents were under low category.

Table - 6 : Distribution of respondents according to their participation in Mass Media

S.N.	Mass Media Participation	Frequency	Percentage
1.	Low (below 6)	22	18.3
2.	Medium (Between 6-9)	51	42.5
3.	High (Above 9)	47	39.2
	Total	120	100.00

Thus, it was concluded that the maximum respondents were from medium category. The findings of the present study were in accordance with the findings of Biswas *et al.* (2008).

Constraints faced by farmers:

Farmers are expected to spend more time with their family members and have domestic work load. In rural areas farmers have little time to spend away from home. Problem in this study has been operationalized as the difficulties faced by the respondent in attending training programme about mushroom cultivation and adoption of mushroom cultivation .Findings related to this aspect of the study have been presented in table 7.

The Table 7 clarified the problems in attending the training programme. The most significant constraint reported was the heavy domestic workload with 80.8% of farmers citing this as a major issue. Other important constraints included the distance problem, as the KVK (Krishi Vigyan Kendra) was far from the village (75.8%), and the lack of freedom of movement for farmers (69.2%). Additionally, farmers reported that they faced challenges related to their ability to spend time away from home (58.3%), a lack of proper facilities

Table - 7 : Problems in attending training programmes .

S.No	Constraints	Frequency	Percentage
1.	Heavy domestic workload.	97	80.8
2.	Less freedom of movement to Farmers.	83	69.2
3.	Little time available to spend on home.	70	58.3
4.	Distance problem (KVK is far away from the village)	91	75.8
5.	Lack of monetary incentives.	53	44.2
6.	Lack of practical experience on mushroom cultivation	63	52.5
7.	Problem in long stay at KVK.	77	64.2
8.	Lack of interest among farmers.	62	51.6
9.	Can't properly understand the technique of mushroom cultivation.	58	48.3
11.	Lack of knowledge about mushroom cultivation.	54	45.0
12.	Lack of proper Facilities at KVK.	64	53.3
13.	Lack of planned workshop for mushroom cultivation	58	48.3
14.	Don't have support of their family.	32	26.6

at the KVK (53.3%), and a lack of practical experience with mushroom cultivation (52.5%). Lack of interest among farmers (51.6%) and the lack of planned workshops for mushroom cultivation (48.3%) were also cited as barriers to adoption. Further analysis revealed that a lack of understanding of the techniques involved in mushroom cultivation (48.3%), a lack of knowledge about mushroom cultivation (45.0%), and a lack of monetary incentives (44.2%) were additional constraints. Finally, some farmers reported that they did not receive support from their family (26.6%) in their efforts to adopt mushroom cultivation.

Suggestions to overcome the constraints faced by mushroom growers in cultivation of mushroom:

Suggestions confronted by mushroom growers in respect of improving their position for further development may be considered as feedback to overcome the constraints. Suggestions were presented in Table 8.

As per the frequency level of the suggestion expressed by the mushroom growers was; there should proper knowledge about mushroom cultivation (rank 1st) followed by proper marketing of the product (rank 2nd), cooperation with mushroom growers (rank 3rd). Continuous

Table - 8 : Suggestions expressed by rural women regarding participation in agricultural activities

S.No	Suggestion	Frequency	Percentage	Rank
1.	There should be family support and guidance.	56	46.6	7
2.	There should be cooperation between family members	61	50.8	5
3.	Proper knowledge about mushroom cultivation	80	66.6	1
4.	Proper marketing of the product	77	64.2	2
5.	Continuous participation in mushroom cultivation training	62	51.6	4
6.	Proper works training should be given	56	46.6	8
7.	There should be proper inspiration about mushroom cultivation	60	50.0	6
8.	Cooperation with mushroom growers	74	61.6	3

participation in mushroom cultivation training(rank 4th).,There should be cooperation between family members(rank 5th) ,There should be proper inspiration about mushroom cultivation(rank 6th). ,There should be family support and guidance(rank 7th) and Proper works training should be given(rank 8th), Pant, Hemlata and Pandey G., (2011), Pant, Hemlata and Pandey, G., (2011), Pant, Hemlata and Pandey, G., (2011), Pant, Hemlata and Pandey, G. (2014).

CONCLUSION

In conclusion, the study highlights the various constraints that farmers face in adopting mushroom production technology in Nalanda, Bihar. Heavy domestic workload, less freedom of movement, and little time available to spend away from home were some of the major constraints faced by mushroom growers. The distance problem as KVK was far from the village, lack of monetary incentives, and lack of practical experience on mushroom cultivation were also significant factors. The long stay in KVK, lack of interest among farmers, and inability to understand the technique of mushroom cultivation were found to increase the difficulties in training programs and of mushroom cultivation. Moreover, the lack of knowledge about mushroom cultivation, lack of proper facilities at KVK, lack of planned workshop for mushroom cultivation and no support from family were also found to affect the adoption of mushroom cultivation. It is important to address these constraints and provide farmers with the necessary resources and support to encourage the adoption of mushroom production technology.

REFERENCES

1. Anchule, M.M. (2000). Critical analysis of technological gap in use of pulse production technology, Ph.D. Thesis, MAU, Parabhani (M.S.).
2. Biswas, S; Datta, M., Sarkar, R. and Sharma, R. (2012). Mushroom as Component of Farming System for Nutrition and Livelihood Improvement in Dhalai. Tripura ICAR Research Complex for NEH Region Tripura Centre. Bulletin 12: 23-29.
3. Biswas, S; Sarkar, A and Goswami, A. (2008). Impact of KVK Training on Advance Dairy Farming Practices (ADFPS) in changing Knowledge and Attitude of Prani-Bandhu. *Journal of Dairying, Foods and Home Science*, 27(1):43-46.
4. Patil, B. V. (2008). Agricultural Indebtedness: Crisis and Revival. *Economics and political weekly*, 43(5):4.
5. Thoke, N. J. and Gunjal, S. (2010). Adoption behavior of farmers in chickpea production technology Department of Agricultural Extension, Krishi Vigyan Kendra, Yashwantrao Chavan Maharashtra Open University, Nashik (M.S.), India. *Agriculture Update*. 5(3/4): 352-355.
6. Pant, Hemlata and Pandey G., (2011): evaluation of different supplements and wheat straw for maximum production of oyster mushroom (P. Sajor caju). *Vignana Parishad Anushandhan Patrika*, vol. 54, No.1: 5-10, [ISSN-0505-5806]
7. Pant, Hemlata and Pandey, G., (2011): influence of wheat straw, paddy straw and saw dust singly and in combination for the maximum production of oyster mushroom (P. sajor caju). *Annl. PI Proc. Sci.* 20 (1) , [ISSN-0971-3571 , impact factor -4-20 and reviewed Journal]. Page No. 259.
8. Pant, Hemlata and Pandey, G., (2011): Effect of different concentration of spent matter (wheat straw) of oyster mushroom (P. Sajor caju) on the growth of tomato plant (*Lycopersicon esculentum*). *J Env. Bio-sci.* Vol. 25 (2): 1-2, [ISSN-0973-6913 (Print) and ISSN 0976-3384 (on line and reviewed Journal)]. Impact factor-4.20.
35. Pant, Hemlata and Pandey, G. (2014): Use of different supplements for the production of oyster mushroom (P. sajor caju) Fr. *Singer. J.Env. Bio. Sci.* Vol. 28(1): 99-100. (ISSN 0976-3384, NASS RATING-4.20).

IMPACT OF DIFFERENT PHOSPHATE SUBSTRATES AND THEIR CONCENTRATION VARIATION OVER PHOSPHATE UPTAKE IN THE CYANOBACTERIUM AULOSIRA FERTILISSIMA

Meenakshee Pandey*, Shalini Singh and Shweta Mishra

*Department of Botany, KNIPSS, Sultanpur, (U.P.), India

E-Mail: 41drmeenakshee@gmail.com

Received : 10.09.2022

ABSTRACT

Accepted : 12.11.2022

The importance of the research's subject matter "Impact of different Phosphate substrates and their concentration variation over Phosphate uptake in the cyanobacterium *Aulosira fertilissima*" remains in the fact that all the N₂-fixing cyanobacteria demonstrated high potential for phosphate acquisition and its accumulation in the form of poly P granules which in turn may be used for phosphate harvesting from fertiliser effluents and eutrophic aquatic ecosystems. The P-uptake behaviour pattern was examined and observed in the test organism in the presence of several P (Inorganic-K₂HPO₄, Na₂HPO₄, and Organic-pNPP) substrates at 30 to 150 minutes of incubation. The data shows that the rate of P uptake is nearly 3.7 times higher in inorganic P (rate of K uptake is 265% more than Na which is 203%) when compared to organic P, even though the latter can serve as an alternative source of P in the absence of inorganic P. However, in both (inorganic & organic) cases within 30 minutes nearly 50 to 70% of substrate has been taken. Under varied K₂HPO₄ supplementation concentrations (1, 5, and 10 mg l⁻¹) in BG-11 growth media, the ability of test organisms to uptake phosphorus was also characterised. Incubation begins with a highly rapid rate of absorption, with 35, 43, and 70% of external P entering cells after 30 minutes, but only 1, 5, and 10% doing so within the following 30 minutes. The work's significance still rests on the fact that P is one of the fundamental constraints on cyanobacterial dominance in natural ecosystems. Throughout the course of the experiment, it became abundantly evident that P has a significant influence on the metabolism of the test organism (*Aulosira fertilissima*). This organism has been discovered to be a powerful biofertilizer with effective P metabolic processes and the capacity to alter its surrounding ecology.

Keywords : *Aulosira fertilissima*, cyanobacteria, various p-substrates, paddy-fields, p-uptake.

INTRODUCTION

Phosphate is essential for development and metabolic activity in all types of life, including cyanobacteria, and is engaged in a number of geochemical processes that are critical in mats. Phosphorus typically only makes up 0.6% of the dry

mass of cyanobacteria cells, however they can occasionally store phosphate as polyphosphate (Stal, 2000).

Orthophosphate (H₃PO₄), the most prevalent form of inorganic phosphorus, is absorbed by cyanobacteria. Elemental concentrations of Ca²⁺,

Mg^{2+} , Fe^{2+} , Fe^{3+} , and Al^{3+} affect how soluble orthophosphate is. Orthophosphate solubility in seawater is primarily regulated by Ca^{2+} , which at an appropriate pH (7.4–8.1) generates the very insoluble hydroxyapatite $\text{Ca}_{10}(\text{PO}_4)_6(\text{OH})_2$; solubility product (1.53X110–112) (Ehrlich, 1996). Additionally, phosphate can also form an insoluble precipitate with the ferric ion ($\text{FePO}_4 \cdot 2\text{H}_2\text{O}$) strengite; solubility product 1.35X10-18 (Stal, 2000).

Microbial activity may allow phosphate to be released from these insoluble minerals. In a distinct genomic island, numerous genes were shown to frequently be present in low-P sites while absent in high-P areas (Martiny et al. 2009). In conclusion, this research shows that *Prochlorococcus*' genome and oceanic environmental circumstances are related.

Two diazotrophic cyanobacteria, *Westiellopsis prolifica* and *Anabaena variabilis*, were tested for their capacity to solubilize Mussorie rock phosphate (MRP) and extracellularly insoluble tricalcium phosphate (TCP). The two strains responded differently to the phosphorus employed in the insoluble forms. According to Yandigeri et al. (2011), TCP (20 mg P l^{-1}) was the greatest source of insoluble P as opposed to MRP or K_2HPO_4 and *A. variabilis*, one of the two cyanobacteria, performed best in terms of P-solubilization and nitrogen fixation.

In phosphorus-limited settings, cellular phosphorous concentrations decrease when the growth rate is reduced, but the potential for phosphorus intake increases. Due to the substantial formation of polyphosphate reserves caused by a pulse of phosphorus delivered to cells with low phosphorus levels (the polyphosphate "overplus" phenomenon), cellular phosphorus levels can exceed those that occur under steady state maximum growth rates (Allen, 1984; Riegman & Mur, 1984). While using a variety of organic phosphates as a substrate, *Calothrix brevissima* and its thermo-tolerant mutant's phosphomonoesterase activity

showed a slight fluctuation in temperature and pH optima, indicating the reliance of temperature and pH optima on the substrate. However, cyanobacterial strains cultivated at their respective temperature and pH maxima differentiated spores less frequently along with early spore commencement. Even though it wasn't apparent until 30°C, the growth of the thermo-tolerant strain (tr) isolated throughout the experiment peaked at 50°C. A sort of thermo-tolerant strain that has been isolated has a lot of potential for use as a biofertilizer in rice farming, particularly in tropical regions (Pandey & Tiwari, 2004).

The uptake rates of phosphate and ATP were less tightly connected in heterotrophic bacteria and *Prochlorococcus* than in *Synechococcus*, and phosphate seemed to be preferred over ATP in all three groups. In comparison to other microbial groupings, *Synechococcus* absorbed both substances at substantially higher rates. In the Sargasso Sea, cyanobacteria compete successfully with other organisms for phosphate and ATP (Michelou et al. 2011).

In order to grow more when they reach the surface, filamentous cyanobacteria can obtain phosphate in deeper strata. They did exist there, albeit in very small numbers. Only 0.3–10% of the biomass in the surface layer at 30 m deep was made up of cyanobacteria. Therefore, compared to other sources, this contribution to bloom development could not be as significant (Nausch et al. 2011).

In microbial cells, inorganic polyphosphate (polyP) plays a significant role in increasing cell resistance to unfavorable environmental conditions and in regulating different biochemical processes. polyP is a polyfunctional compound. The most important of its functions are the following: phosphate and energy reservation, cat-ion sequestration and storage, membrane channel formation, participation in phosphate transport, involvement in cell envelope formation and function, gene activity control, regulation of enzyme activities, and a vital role in stress response

and stationary-phase adaptation. The functions of polyP have changed greatly during the evolution of living organisms. In prokaryotes, the most important functions are as an energy source and a phosphate reserve. In eukaryotic microorganisms, the regulatory functions predominate. Therefore, a great difference is observed between prokaryotes and eukaryotes in their polyP-metabolizing enzymes (Kulaev & Kulakovskay, 2000).

Phosphorus potentially limits the growth and productivity of the unicellular cyanobacterium *Synechococcus* in many oligotrophic oceans. Both dissolved inorganic phosphorus (DIP) and dissolved organic phosphorus (DOP) could be important P sources in these regimes (Fu et.al., 2006).

Phosphorus availability is one of the key factors implicated in controlling cyanobacterial growth (Armstrong, 1999). Phosphorous (P) starved cells of the cyanobacterium *Anabaena oryzae* showed higher phosphate uptake rates than P-sufficient cells. The P-uptake obeyed saturation kinetics. The K_m value for P-deficient cells was lower ($54.34 \mu\text{M}$) than P-sufficient cells ($82.64 \mu\text{M}$) while V_{max} was higher in P-deficient and lower in P-sufficient cells (Singh et.al. 2007).

In the Baltic Sea, surplus phosphorus after the spring bloom or phosphorus intake from deeper levels, such as by upwelling, are nutritional sources for the growth of filamentous cyanobacteria (Nausch et al. 2011).

In *Prochlorococcus* uptake of PO_4 depends on the high-affinity P uptake system composed of a periplasmic-binding protein, PstS, and associated membrane-bound ABC transporter, PstCAB (Scanlan et al., 2009). Expression of these four genes is coordinately regulated (Martiny et al., 2006). When Pi is depleted, cells attempt to scavenge P from organic sources by upregulating *phoA* to produce alkaline phosphatase (APase), which can cleave P from organic compounds such as b-glycerophosphate or glucose-6-phosphate (Krumhardt et al., 2013).

Cyanobacterial blooms are generally

explained by nutrient uptake kinetic constants that may confer competitive capabilities, like high nutrient-uptake rate, affinity, and storage capacity. However, cyanobacteria are capable of flexible physiological responses to environmental nutrient fluctuations through adaptation of their phosphate uptake properties. Growth optimization is possible if the physiological reaction time of cyanobacteria (tR) matches the duration of nutrient availability (Aubriot & Bonilla, 2012).

Synechocystis cells can survive under Pi-limiting conditions following initial growth in BG-11 medium. The uptake of Pi in *Synechocystis* 6803 is accomplished mainly by Pst1 despite its lower affinity for Pi than that of Pst2. The expression of Pst2 might be useful when cells encounter low Pi environments. Pi uptake is stimulated by alkaline pH as well as by ionic solute such as NaCl whereas it is inhibited by non-ionic solute (sorbitol) generating osmotic stress (Burut-Archanai et.al., 2011).

All living things require phosphorus, a vital mineral nutrient, for a variety of processes, including the production of DNA, RNA, and nucleotides like ATP, as well as the functional control of proteins through phosphorylation. Although it is typically present in natural surroundings at nanomolar levels, inorganic phosphate (Pi), the only form of phosphorus that can be utilised directly by organisms, is frequently limited (Hudson et al., 2000).

The fact that there were no significant differences in Pi uptake at pH 7 and 10 suggested that the Pi uptake system in *Synechocystis* 6803 can recognize both H_2PO_4^- and HPO_4^{2-} . The ability of *Synechocystis* 6803 to bind two different Pi species is advantageous to its survival especially under fluctuating external pH and low Pi availability (Lindsay, 1979).

The increased Pi uptake activity by NaCl is ascribed to an ionic rather than an osmotic effect since an osmotic stress of the same strength achieved with a non-ionic sorbitol caused a reduction in Pi uptake. It is possible that the presence of Na^+ might facilitate

the uptake of Pi, as in *E. coli* where it is transported as neutral metal phosphate (van Veen et.al.,1994). The driving force for the uptake of Pi in *Synechocystis* 6803 is likely to be ATP generated by ion gradient or ion gradient itself. Indeed, the effect of the inhibitors tested on this uptake support this hypothesis. The fact that Pi uptake is Na⁺-stimulated and that the uptake is favorable at alkaline pH can support this contention (van Veen et.al.,1994).

Florida Bay, a shallow, seagrass-dominated bay in southern Florida, USA, absorbs a lot of nutrients and has recently seen microalgal blooms and seagrass losses. Although the effects of organic nutrients, specifically dissolved organic nitrogen (DON) and organic phosphorus (DOP), on microbial processes in the bay are unknown, inorganic nutrient inputs have been widely studied. There are significant management repercussions from the discovery that DON and DOP may contribute differently to the growth of cyanobacteria and other microorganisms in Florida Bay (Glibert, 2004).

MATERIALS AND METHODS

The diazotrophic heterocystous cyanobacterium *Aulosira fertilissima* Ghose was used as experimental materials and it was obtained from paddy fields in epiphytic form.. The present study was conducted in rice-grown areas associated with Lucknow zone which phytographically belongs to Indo-Gangetic Plain. The strain was raised in cultures using standard microbiological culture techniques (Norris & Ribbons, 1968). The culture media, modified Chu-10 (Safferman & Morris, 1964) and BG-11 (Rippka et.al. 1979) were used for routine transfer and maintenance of cyanobacterial cultures. The pH of the medium was adjusted to 7.6 by HEPES (0.2 M) buffer.

Assay of Phosphate Uptake:

The P-acquisition in the N₂-fixing strains was determined on the basis of depletion of phosphate from the medium. The exponential phase culture cells were harvested through centrifugation

(10000 x g: 10 min.) washed with sterilized diluted culture (BG-11) medium and homogenized with glass beads. The cyanobacteria cells in the form of homogeneous suspension were inoculated in P-deficient BG-11 medium with 1, 5, and 10 mg P l⁻¹ substrate (K₂HPO₄) concentrations. The level of phosphate in the medium was determined at 30 min to 150 min. of incubation. The rate of P-acquisition per unit of protein cell biomass was determined in the medium. The P-uptake in other P substrates like Na₂HPO₄ and pNPP was detected at 10 mg P l⁻¹ concentration in a similar experiment described as above.

Cells grown in BG-11 medium for 3 days were washed twice by centrifugation and resuspension in Pi-limiting BG-11 medium. The washed cells were subsequently grown in either BG-11 or Pi-limiting BG-11 medium for 24 h before being washed twice by centrifugation and resuspension in Pi-free buffer to an optical density at 730 nm of 0.3. The uptake experiment was initiated by the addition of K₂HPO₄ solution at room temperature. At different time intervals, aliquots were withdrawn, filtered through a 0.45 µm membrane filter and the remaining Pi in the filtrate was determined by the colorimetric method (Katewa & Katyare, 2003). The statistical analysis of the data were performed using analysis of variance (ANOVA) as described by Underwood (1997).

RESULTS AND DISCUSSTION

The *Aulosira fertilissima* is a dominant cyanobacteria of rice field appeared throughout the crop season where it can be used as a potent biofertilizer. The *A. fertilissima* was incubated under different growth conditions in terms of P supplementation in separate experimental set up.

Influence of different P-Substrates& their variable levels (concentration) over Phosphate uptake:

Cyanobacteria are frequently prominent members of undesirable blooms in natural waters, in which phosphate availability is a common limiting factor. The phosphorus contents of acid soluble pools, lipid, ribonucleic acid, and insoluble polyphosphate were

lowered in organisms in proportion to the reduction in growth rate under phosphate-limited but not in nitrate-limited continuous culture. Phosphorus in these cell fractions was lost proportionately during progressive phosphate starvation of batch cultures. In the previous experiments, a differential response of diazotrophic cyanobacterial strain *Aulosira fertilissima* in terms of growth was observed in relation to the different P and N- sources. The growth in the presence of phosphate depends upon the cyanobacterial capacity to utilize the phosphate. The utilization occurs in a two-step process, uptake of the nutrient and its further metabolism. The potential of *A. fertilissima* to take up phosphorus by phosphate starved cells under different P substrates was investigated during present course of work.

A. Phosphate uptake behavior under different P-substrates:

It is widely accepted that phosphorus is the main nutrient that controls the development of natural populations of cyanobacteria since, depending on its concentration; it can act as a growth limiting factor or, on the other hand, promote blooms in which N₂ fixing cyanobacteria prevent the population from becoming N-deficient. Orthophosphate is believed to be the form of phosphorus commonly taken up by autotrophs. Under normal phosphate supply, its uptake by cyanobacteria is an apparent hyperbolic function of the external phosphate concentration but generally phosphate uptake is modified under conditions of P deficiency. Although the cyanobacteria potential to use organic sources of P cannot be generalized, the utilization of phosphate monoesters has been shown on different occasions. The ability is due to the fact that the cyanobacteria contain extracellular phosphatases that release the orthophosphate. Cell-bound alkaline phosphatases are typical of phytoplankton and seem to respond to P-deficiency since they are adaptative enzymes depending on phosphate concentration.

Keeping in view of the above the P-uptake was investigated in test organisms in presence of

different P-substrates (K₂HPO₄, Na₂HPO₄, and pNPP) and the result has been presented in the Figure 1. Cyanobacterial strains starved for phosphate for 72 h were harvested, washed and incubated in different P-substrates supplemented BG-11 growth medium and P-uptake was determined at 30 min to 150 min of incubation.

It is quite evident from the findings obtained (Fig. 1) that inorganic P is an essential requirement of an organism in comparison to organic P, however the latter can serve as a provisional source in absence of available orthophosphate. P uptake rate is nearly 3.7 times higher in inorganic P (K is preferred 265% more over Na 203% as nutrient) when compared to organic P; however in both cases within 30 minutes nearly 50 to 70 % of substrate has been taken by the organism.

B. Substrate concentration dependent Phosphate uptake pattern:

Keeping in view of the above the P-uptake was further investigated and characterized by taking K₂HPO₄ as substrate and varying its concentration in test organism and the result has been presented in the Figure 2. Cyanobacterial strains starved for phosphate for 72 h were harvested, washed and incubated in different concentration supplemented BG-11 growth medium.

P-uptake was determined at 30 min to 150 min of incubation under 1, 5 and 10 mg l⁻¹ and the result reveals that the P-uptake phenomenon determined in test organism is a concentration dependent mechanism. The rate of uptake is very fast within 30 minutes and 35, 43, and 70% of external P (1mg, 5mg, and 10mg P l⁻¹, respectively) has been transported intracellular, however this rate gradually slowed down.

Cyanobacterial blooms caused by eutrophication in Lake Taihu have led to ecological threats to freshwater ecosystems (Peng et. al., 2020). A pilot scale experiment was implemented to investigate the relationship between cyanobacteria and other aquatic plants and animals in simulated eutrophic ecosystems under different phosphorus

(P) regimes. The results of this study showed that cyanobacteria had two characteristics favorable for bloom formation in eutrophic ecosystems. One is the nutrient absorption. The presence of alkaline phosphatase was beneficial for algal cells in nutrition absorption under low P concentration. The other is the release of microcystin.

Many photosynthetic microorganisms, including cyanobacteria, evolved in nutrient-poor environments or environments with varying nutrient concentrations. As a result, they acquired a broad array of acclimations to scarce and/or fluctuating nutrient availability, including the capability for luxury uptake. Although this phenomenon has been known for quite a long time (Ketchum, 1939), many aspects of LPU remain largely unknown to date.

Phosphorus (P) is an important nutrient central to storing and the exchange of energy and information in the cell (Dyhrman, 2016). At the same time, the availability of P in many habitats dwelled by cyanobacteria is scarce and/or fluctuating. Cyanobacteria developed a broad array of acclimations to cope with P shortage. One of the most widespread is the capability of taking up P in large excess of the current metabolic demand, termed “luxury P uptake” (LPU) (Ketchum, 1939). The ability of the P-deprived culture to accumulate

excessive amounts of inorganic polyphosphate (PolyP) after refeeding with P_i is known as “hypercompensation” or “phosphate overplus” (Kulaev, 2004).

To cope with fluctuating phosphorus (P) availability, cyanobacteria developed diverse acclimations, including luxury P uptake (LPU)—taking up P in excess of the current metabolic demand (Solovchenko et. al., 2020). LPU is underexplored, despite its importance for nutrient-driven rearrangements in aquatic ecosystems. Solovchenko et. al. (2020) studied the LPU after the refeeding of P-deprived cyanobacterium *Nostoc* sp. PCC 7118 with inorganic phosphate (P_i), including the kinetics of P_i uptake, turnover of polyphosphate, cell ultrastructure, and gene expression. The P-deprived cells deployed acclimations to P shortage (reduction of photosynthetic apparatus and mobilization of cell P reserves). The P-starved cells capable of LPU exhibited a biphasic kinetic of the P_i uptake and polyphosphate formation. The first (fast) phase (1–2 h after P_i refeeding) occurred independently of light and temperature. It was accompanied by a transient accumulation of polyphosphate, still upregulated genes encoding high-affinity P_i transporters, and an ATP-dependent polyphosphate kinase. During the




Figure - 1 : Phosphate uptake behavior of *Aulosira fertilissima* under different P- substrates.

Anova: Single Factor						
SUMMARY						
Groups	Count	Sum	Average	Variance		
Column 1	6	12	2	1.572		
Column 2	6	17.9	2.983333333	3.045666667		
Column 3	6	30.5	5.083333333	6.833666667		
ANOVA						
Source of Variation	SS	df	MS	F	P-value	F crit
Between Groups	29.76777778	2	14.88388889	3.899254817	0.0432898	3.682320344
Within Groups	57.25666667	15	3.817111111			
Total	87.02444444	17				

- The group comparison test (one way ANOVA) for three different samples i.e., 1mg PI^{-1} , 5 mg PI^{-1} and 10 mg PI^{-1} supplemented growth conditions for *Aulosira fertilissima* cells were performed for determining the possibility of variation among the group sample means is greater than expected by chance or not i.e., the null hypothesis is being tested.
- The F test is performed by taking the ratios of among MS and within MS.
- $F_{\text{critical/Expected}} = 3.682320344$ and
- $F_{\text{observed}} = 3.899254817$ at alpha (α) = 0.05 and df = 2, 15.
- Here, the calculated F-value (3.89) is larger than F_{critical} (3.68) for $\alpha = 0.05$.
- Thus, the test is significant at $\alpha = 5\% \rightarrow$ Not all mean samples are the same; the substrate concentration acts as an influential factor for cyanobacterial growth and phosphate uptake behavior.
- Since $F_{\text{observed}} > F_{\text{Expected}}$ so null hypothesis is rejected.
- It can be concluded that there is significant inequality among the three test samples (i.e., one or more sample is significantly different from the othersamples).




Figure - 2 : Substrate concentration dependent Phosphate uptake behavior of *Aulosira fertilissima*.

Anova: Single Factor						
SUMMARY						
Groups	Count	Sum	Average	Variance		
Column 1	6	12	2	1.572		
Column 2	6	17.9	2.983333333	3.045666667		
Column 3	6	30.5	5.083333333	6.833666667		
ANOVA						
Source of Variation	SS	df	MS	F	P-value	F crit
Between Groups	29.76777778	2	14.88388889	3.899254817	0.0432898	3.682320344
Within Groups	57.25666667	15	3.817111111			
Total	87.02444444	17				

- The group comparison test (one way ANOVA) for three different samples i.e., 1 mg PI^{-1} , 5 mg PI^{-1} and 10 mg PI^{-1} supplemented growth conditions for *Aulosirafertilissima* cells were performed for determining the possibility of variation among the group sample means is greater than expected by chance or not i.e., the null hypothesis is being tested.
- The F test is performed by taking the ratios of among MS and within MS.
- $F_{\text{critical/Expected}} = 3.682320344$ and
- $F_{\text{observed}} = 3.899254817$ at $\alpha (\alpha) = 0.05$ and $df = 2, 15$.
- Here, the calculated F-value (3.89) is larger than F_{critical} (3.68) for $\alpha = 0.05$.
- Thus, the test is significant at $\alpha = 5\% \rightarrow$ Not all mean samples are the same; the substrate concentration acts as an influential factor for cyanobacterial growth and phosphate uptake behavior.
- Since $F_{\text{observed}} > F_{\text{Expected}}$ so null hypothesis is rejected.
- It can be concluded that there is significant inequality among the three test samples (i.e., one or more sample is significantly different from the others samples).

second (slow) phase, recovery from P starvation was accompanied by the downregulation of these genes. The study revealed no specific acclimation to ample P conditions in *Nostoc* sp. PCC 7118.

CONCLUSION

Agrochemical companies and farmers are interested in cyanobacteria because of their abilities as biostimulants and biofertilizers. Crop production may use cyanobacterial biostimulants and biofertilizers to promote agricultural sustainability. Hence, cyanobacteria are employed in agriculture in different applications, such as amendment, foliar application, and seed priming. These may be used as a substitute for synthetic fertilisers, crop protection agents, and plant growth regulators, or in combination with them to produce a variety of benefits, including improved roots, higher crop

yields and quality, and tolerance to salt and drought. This study is aimed at reporting the uptake potential of *Aulosira fertilissima* under different inorganic and organic P-substrates and under different concentrations of inorganic P-substrate (K_2HPO_4). Cyanobacteria showed better acquisition potential under inorganic P-substrates in comparison to organic P- substrates. Hence, it can be said that the organic P-substrates would be used as an alternated source of P in the case of inorganic P deficiency.

ACKNOWLEDGEMENTS

The authors thank Head, Department of Botany, KNIPSS, Sultanpur for providing the facilities. Authors are grateful to colleagues and staff members for encouragements.

REFERENCES

1. Allen M.M. (1984): Cyanobacterial cell

- inclusions. *Annu. Rev. Microbiol.*, 38:1-25.
2. Armstrong, R.A. (1999). An optimization-based model of iron light-ammonium co-limitation of nitrate uptake and phytoplankton growth. *Limnol. Oceanogr.*, 44: 1436–1446.
3. Aubriot L., Bonilla S. (2012): Rapid regulation of phosphate uptake in freshwater cyanobacterial blooms. *Aquatic Microbial Ecology*, 67: 251–263. doi: 10.3354/ame01596.
4. Burut-Archanai S., Eaton-Rye J.J., and Incharoensakdi A. (2011): Na⁺ -stimulated phosphate uptake system in *Synechocystis* sp. PCC 6803 with Pst1 as a main transporter. *BMC Microbiology*, 11:225-231.
5. Dyhrman S.T. (2016): *The Physiology of Microalgae*. Springer; Berlin/Heidelberg, Germany. Nutrients and their acquisition: Phosphorus physiology in microalgae; pp. 155–183.
6. Fu F-X., Zhang Y., Feng Y., Hutchins D.A. (2006): Phosphate and ATP uptake and growth kinetics in axenic cultures of the cyanobacterium *Synechococcus* CCMP 1334. *Eur. J. Phycol.*, 41: 15–28.
7. Glibert P.M., Heil C.A., Hollander D., Revilla M., Hoare A., Alexander J., Murasko S. (2004): Evidence for dissolved organic nitrogen and phosphorus uptake during a cyanobacterial bloom in Florida Bay. *Marine Ecology Progress Series*, 280: 73–83.
8. Hudson JJ, Taylor WD, Schindler DW (2000): Phosphate concentrations in lakes. *Nature*, 406: 54-56. 10.1038/35017531.
9. Katewa SD, Katyare SS (2003): A simplified method for inorganic phosphate determination analysis in enzyme assays. *Anal Biochem.*, 323: 180-187. 10.1016/j.ab.2003.08.024.
10. Ketchum B.H. (1939): The absorption of phosphate and nitrate by illuminated cultures of *Nitzschia closterium*. *Am. J. Bot.*, 26:399–407. doi: 10.1002/j.1537-2197.1939.tb09293.x.
11. Krumhardt, K., Callnan, K., Roache-Johnson, K., Swett, T., Robinson, D., Nahas Reistetter, E. (2013): Effects of phosphorus starvation versus limitation on the marine cyanobacterium *Prochlorococcus* MED4 I: uptake physiology. *Environ Microbiol.* doi: 10.1111/1462-2920.12079.
12. Kulaev I., Vagabov I., Kulakovskaya T. (2004): *The Biochemistry of Inorganic Polyphosphates*. 2nd ed. John Wiley & Sons, Ltd; Chichester, UK.
13. Kulaev, I., Kulakovskay T.V. (2000): Polyphosphate and Phosphate Pump. *Annual Review of Microbiology* 54(1):709-34. DOI:10.1146/annurev.micro.54.1.709
14. Lindsay WL (1979): Chemical equilibria in soils. Wiley-Interscience Publ., New York.
15. Martiny A.C., Huang V. and Weizhong L.F. (2009): Occurrence of phosphate acquisition genes in *Prochlorococcus* cells from different ocean regions. *Environmental Microbiology*, 11(6):1340-1347.
16. Martiny, A.C., Coleman, M.L., and Chisholm, S.W. (2006): Phosphate acquisition genes in *Prochlorococcus* ecotypes: evidence for genome-wide adaptation. *Proc Natl Acad Sci, USA* 103: 12552–12557.
17. Michelou V.K., Michael L.W. and David K.L. (2011); Phosphate and adenosine-5'-triphosphate uptake by cyanobacteria and heterotrophic bacteria in the Sargasso Sea. *Limnol. Oceanogr.*, 56(I):323-332.
18. Nausch M., Nausch G., Mohrholtz V., Siegel H. and Wasmund N. (2011): Is growth of filamentous cyanobacteria supported by phosphate uptake below the thermocline?

Estuarine, Coastal and Shelf Science, 99:50-60.DOI: 10.1016/j.ecss.2011.12.011.

19. Norris J.R. and Ribbons D.W. (1968): Methods in Microbiology. Vol.3. Academic Press, London and New York.

20. Pandey M. and Tiwari D.N. (2004): A Correlation Matrix of Alkaline Phosphatase and Sporulation in Diazotrophic Cyanobacteria and its Thermo-tolerant Mutant. Polish Journal of Microbiology, 53(4):257-265.

21. Peng G., Li Q., Zhang H., Luo X., Zhang W., Zheng Z., Luo X. (2020): Effects of Cyanobacteria on Phosphorus Cycling and Other Aquatic Organisms in Simulated Eutrophic Ecosystems. Water, 12, 2265; doi:10.3390/w12082265.

22. Riegman R. and Mur L.C. (1984): Regulation of phosphate uptake kinetics in Oscillatoria agardhii. Arch. Fur Microbiol., 139:28-32.

23. Rippka R., Deruelles J., Waterbury J.B., Herdman M. and Stainer R.Y. (1979): Generic assignment, strain histories and properties of pure cultures of cyanobacteria. J. Gen. Microbiol., 111:1-61.

24. Safferman R.S. and Morris M.E. (1964): Growth characteristics of the blue green algal virus LPP-1. J. Bacteriol., 88:771-775.

25. Scanlan, D.J., Ostrowski, M., Mazard, S., Dufresne, A., Garc zarek, L., and Hess, W.R. (2009): Ecological genomics of marine picocyanobacteria. Microbiol. Mol. Biol. Rev. 73:249–299.

26. Singh S.K., Pandey V., Pandey K.D. (2007): Phosphate uptake kinetics and its regulation in N2- fixing cyanobacterium Anabaena oryzae Fritsch under salt stress. African Journal of Biotechnology Vol. 6 (20), pp. 2363-2368.

27. Solovchenko A, Gorelova O, Karpova O, SelyakhI, Semenova L, Chivkunova O,

Baulina O, Vinogradova E, Pugacheva T, Scherbakov P, Vasilieva S, Lukyanov A, Lobakova E (2020): Phosphorus Feast and Famine in Cyanobacteria: Is Luxury Uptake of the Nutrient Just a Consequence of Acclimation to Its Shortage? Cells, 9(9):1933. doi: 10.3390/cells9091933. PMID: 32825634; PMCID: PMC7564538.

28. Stal L.J. (2000): Cyanobacteria Mats and Stromatolites. In: B.A. Whitton and M. Potts (eds) The Ecology of Cyanobacteria, pp.61-120, Kluwer Academic Publishers, The Netherlands.

29. Underwood A.J. (1997): Their logical design and interpretation using analysis of variance. Experiments in Ecology, Cambridge University Press, ISBN 0-521-55329-6.

30. van Veen HW, Abee T, Kortstee GJ, Konings WN, Zehnder AJ (1994): Translocation of metal phosphate via the phosphate inorganic transport system of Escherichia coli. Biochemistry, 33: 1766-1770. 10.1021/bi00173a020.

31. Yandigeri M.S., Meena K.K., Srinivasan R. and Pabbi S. (2011): Effect of mineral phosphate solubilization on BNF by diazotrophic cyanobacteria. Indian Journal of Microbiology, 51(I): 48-53.

ASSESSING THE CHALLENGES HINDERING FARM WOMEN'S EFFECTIVE USE OF MASS MEDIA FOR EMPOWERMENT AND RURAL DEVELOPMENT

Km. Arti, N. K. Mishra and Pradeep Kumar Yadav

Department of Agricultural Extension

T. D. P.G. College, Jaunpur, (U.P.), India

Corresponding email: Prajapatikmarti@gmail.com

Received : 08.01.2023

ABSTRACT

Accepted : 11.02.2023

This study aimed to assess the challenges hindering farm women's effective use of mass media for empowerment and rural development. Through a comprehensive analysis of constraints, including lifestyle factors and mass media utilization barriers, a range of challenges were identified. These challenges encompassed farm women's busy schedules with household work, unreliable electricity due to load shedding, poor economic conditions, limited time for utilizing mass media, inadequate educational opportunities, lack of confidence resulting from societal pressures and customs, and insufficient knowledge and communication facilities. The findings highlight the need for concerted efforts to overcome these constraints and promote the empowerment of farm women, as well as induce a positive change in their attitudes toward social issues. Strategies to address these challenges include providing training and educational opportunities, improving infrastructure, promoting confidence and empowerment, tailoring content and delivery, and fostering collaboration and partnerships among various stakeholders. By addressing these constraints and implementing the recommended strategies, it is possible to create an environment that supports farm women in effectively utilizing mass media for their empowerment and rural development. This, in turn, can contribute to significant social transformations and enhance the overall well-being of farm communities.

Keywords : *Unreliable electricity, load shedding, empowerment, opportunities .*

INTRODUCTION

The mass media is the vehicle that carry messages to a large audience. It is very pervasive in modern life but normally people do not even notice its influence. In liberal democracy, the role of the media can be summarised as a medium of information, entertainment and a medium to educate the people. It is widely accepted in liberal democracy that when the media help to put

information at the disposition of the people, they will be able to formally or informally control the state. The mass media is essential for democracy. By keeping people on top of current issues, the media enables people to participate intelligently in public policy discussion and decision- making. In a democracy the principal role of the media is to act as a check on the state and fearlessly expose abuses of official authorities. This watchdog role is said to

override in importance all other functions of the media and dictate the form in which the media system should be organized. The media is also the vehicles by which people debate the issues and try to persuade each other is different points of view. Even when it provides us entertainment, the mass media is capable of portraying and shaping values that enrich our dialogue on social issues and public policy (Baishakhi Nag, 2011).

The mass media has long been welcomed as the watchdog of society and this tradition bestows upon them the social responsibility to mirror and guide the process of social change. But in contemporary India, mass media under various pressures has become the commercial channels only and failed to reflect the social problems or aspirations of the entire population, especially the problems faced by women in particular. So, the portrayal of women in Indian media is films, television programmes, visual advertisements or newspaper and magazine is becoming an area of great concern to the people having interest in social research and studies. There is an on-going trend

MATERIALS AND METHODS

The present study has been carried out in the Jaunpur district of Uttar Pradesh. Out of 75 district in Uttar Pradesh, Jaunpur district was selected purposively. There are 21 blocks in Jaunpur district. Out of which one block namely Sondhi was selected purposively based on the assumption that the Tilak Dhari Post Graduate College, Jaunpur is well connected center for education and transfer of latest agricultural technology. Hence, this block was selected.

The study was conducted in Sondhi block of Jaunpur district (Uttar Pradesh) during 2021-22. Main purpose of the study is to assess the impact of mass media on life style of farm women. From the selected block, five villages were selected for the present investigation through random sampling method. Thus five villages were selected for the study purpose. The selection of these villages is based on the assumption that these were having

impact of mass media on lifestyle of farm women. From each selected village 24 respondents were randomly selected. Thus the total sample size constituted to 120 respondents. A structured questionnaire was developed to gather quantitative data on the challenges faced by farm women in utilizing mass media. The questionnaire will include closed-ended questions to measure the frequency and severity of challenges.

RESULTS AND DISCUSSION

Social participation:

Social participation of farm women exposes them to various spheres of life as it leads to their empowerment. It is important parameter in social empowerment. The social participation of women respondents had been studied in the present investigation.

It was observed from the table 1 that majority of women respondents 78.3 per cent had occasional social participation, 15.8 per cent women respondents had never and 5.8 per cent women respondents had regular social participation.

Table - 1 : Distribution of the respondents according to their social participation:

S. N.	Category	Frequency	Percentage
1.	Regularly	7	5.8
2.	Occasionally	94	78.3
3.	Never	19	15.8
	Total	120	100.0

It is concluded that majority of respondents were found in the occasional category of social participation.

The same findings were reported by Lohare (2017). That 78.3 per cent respondents has participated in social organization in occasional category.

Attitude towards mass media:

It was observed from the table 2 that majority of women respondents 71.7 per cent agreed

with attitude towards mass media, 18.3 per cent women respondents undecided and 10.00 per cent women respondents had strongly agreed with mass media.

Table - 2 : Distribution of the respondents according to their Attitude towards mass media:

S. N.	Category	Frequency	Percentage
1.	Strong Agree	12	10.0
2.	Agree	86	71.7
3.	Undecided	22	18.3
	Total	120	100.0

These findings were in conformity with the findings of Rambabu *et. al.* (2000). They have reported similar observations where in most of the farm women members were 71.7 per cent agree with attitude towards mass media.

Availability of mass media:

It was observed from the table 3 that majority of women respondents (75 per cent) have availed mass media with difficulty. 13.3 per cent easily available to mass media and 11.6 per cent not available to mass media.

Table - 3 : Distribution of The Respondents according to their availability of mass media:

S. N.	Category	Frequency	Percentage
1.	Easily available	16	13.3
2.	Available with difficulty	90	75.0
3.	Not available	14	11.6
	Total	120	100.0

Due to exposure and viewing to various sources of information women's get influenced to do something new to increase their socio-economic status. From this point of view the media availability has been considered for the present study.

Challenges hindering farm women's in effective use of mass media for empowerment and rural development

Table - 4 : Constraints in effective use of mass media.

Constraints	Percentage	Rank
Too busy in household work	59.33%	I
Improper electricity due to load shedding	58.00%	II
Poor economic condition	55.33%	III
Lack of time for utilizing mass media	50.67%	IV
Inadequate educational opportunities	42.67%	V
Farm women feel inhibited due to lack of confidence, family pressure, and customs etc.	39.33%	VI
Lack of knowledge	34.67%	VII
Poor communication facilities	30.67%	VIII

The constraints analysis revealed from table-4 that the life style and mass media utilization constraints i.e., too busy in household work ranked as the highest concern, with a percentage of 59.33%. Improper electricity due to load shedding was the second-highest constraint, with a percentage of 58.00%. Poor economic condition ranked third, with a percentage of 55.33%. Lack of time for utilizing mass media accounted for 50.67% of the constraints. Inadequate educational opportunities were reported by 42.67% of the participants, while farm women feeling inhibited due to lack of confidence, family pressure, and customs accounted for 39.33%. Lack of knowledge and poor communication facilities were constraints reported by 34.67% and 30.67% of farm women, respectively. These data highlight the multifaceted challenges that farm women face in effectively utilizing mass media for their empowerment and rural development. Addressing these constraints it is crucial to enable farm women to overcome barriers and fully leverage the potential of mass media for their benefit. It could be concluded that the analysis of these constraints emphasizes to have concerted efforts to overcome these so as to increase the empowerment of farm women and to change their attitude towards the social issues.

CONCLUSION

In conclusion, the assessment of challenges hindering farm women's effective use of mass media for empowerment and rural development revealed that a majority of the respondents had low social participation, while a significant proportion

exhibited a positive attitude towards mass media. Television, radio, and newspapers were readily available, but other print media options were lacking. The analysis highlighted constraints related to lifestyle and mass media utilization, such as busy household work, unreliable electricity, poor economic conditions, lack of time, limited educational opportunities, lack of confidence, family pressure, customs, inadequate knowledge, and poor communication facilities. Overcoming these constraints requires concerted efforts to increase farm women's empowerment and transform their attitudes towards social issues. By addressing these challenges, we can enhance their access to and effective use of mass media, fastering their empowerment and contributing to rural development.

REFERENCES

1. Kumar B. T.P. (2010). Time utilization and decision making in horticulture: Antecedent to gender mainstreaming. M.Sc. (Agri.) Thesis (Unpub.), Univ. Agri. Sci., Bengaluru.

2. Kumar, A. Paswan, A. K. Ansari, M. N. and Singh, A. K. (2016). Relationship of socio-

economic profile of the turmeric growing farmers to their training need.*Indian Journal of Extension Education*.52 (2): 97-100.

3. Kumar, P. Kher, S. K. Nain, M.S. and Slathia, P. S. (2018).Socio-economic assessment of farmer in 'kandi' region of Jammu.*Indian Journal of Extension Education*.54 (3): 146-149.

4. Rathod, R. and Trikha, R. N. (2006). Information need and utilization pattern of subscriber of RAU publications. *International J. Ext. Educ.* Dec. 2006: 95-97.

5. Rambabu, P.P., Ramaiah V. and Rao, P.P. (2000). A Scale to measure attitude of Farmers towards information technological knowledge. *Maharashtra J. Ext. Educ Vol.XIX*: 82-85.

6. Ram, D. Mark, R.T. and Chaudhary, K.P. (2014).Mass media exposure of north – Eastern Hill University students.*Journal of communication studies*, vol. XXXII (pp. 11_18).

IMPACT OF KVK TRAINING PROGRAM ON KNOWLEDGE AND ADOPTION OF MUSHROOM PRODUCTION TECHNOLOGY

Anish Kumar, N. K. Mishra and Pradeep Kumar Yadav

Department of Agricultural Extension

T. D. P.G. College, Jaunpur, (U.P.), India

Corresponding email: anish8084kr@gmail.com

Received : 25.12.2022

ABSTRACT

Accepted : 28.01.2023

This study aimed to assess the impact of KVK (Krishi Vigyan Kendra) training programs on the knowledge and adoption of mushroom production technology among farmers in District Nalanda, Bihar. A total of 120 respondents were surveyed with most of them being young, male, unmarried and belonging to the OBC category. The majority of respondents had a graduate education and were engaged in primary occupations and the most common family type was joint. The study found that most respondents had a medium level of knowledge and adoption of mushroom production technology. However after attending KVK training programs there was a significant improvement in the level of knowledge and adoption of mushroom production technology among the respondents. The study concludes that KVK training programs have a positive impact on farmers' knowledge and adoption of mushroom production technology which can lead to increased productivity and income in the agricultural sector.

Keywords : *Adoption, productivity, technology .*

INTRODUCTION

Indian economy is based on the growth of agriculture. After 1964-65 the tremendous production is observed in our country. In present time the production is stagnant in major food crops. There is a need to strengthen extension system which plays a key role in modernising country so that production can be extended. Training is one of the important component of extension system. The directorate of Extension and the Ministry of Agriculture funding to all types of training. The KrishiVigyan Kendra (KVKs) are functioning as per the mandates to transfer the latest agriculture

technology to the farmers' field through organising short and long-term training courses for farmers, farm-women, rural youth and extension workers, laying down the demonstration in the farmers' field with their participation. Dr. Mohan singh Mehta committee recommended the establishment of a network of KrishiVigyan Kendra (KVKs) in the country.

Mushroom growing is one agricultural activity in which farmers can play a vital role without sacrificing their household responsibilities. Krishi Vigyan Kendra, Nalanda is imparting trainings in mushroom cultivation to the farmers,

farm women and rural youth. During 2018-19 to 2020-21, five such vocational training programmes valuate the outcome of these training programmes, a study was conducted to assess the socio - economic profile of the trainee, gain in knowledge and adoption status of the enterprise in mushroom farming among the rural farmers of the Nalanda District were organized regarding mushroom farming in which 120 farmers participated. In order to evaluate the outcome of these training programmes, a study was conducted to assess the socio- economic profile of the trainee, gain in knowledge and adoption status of the enterprise in mushroom farming among the rural farmers of the Nalanda District.

The Oyster mushroom (*Pleurotus* sp.) belonging to class Basidiomycetes and family Agaricaceae is popularly known as Dhingri in India and grows naturally in the temperate and tropical forests. It also grows on decaying organic matter and produces protein rich food. The best growing season is September/October to March/April. It requires a short growth time in comparison to other edible mushrooms. The economic importance of the mushroom lies primarily in its use a food for human consumption. It is rich in vitamin C and B complex and the protein content varies between 1.6 to 2.5 per cent. Farmers were mainly focused on button mushroom than Dhingri. Mushroom cultivation can directly improve livelihoods through economic, nutritional and medicinal contributions. Mushroom is a popular food due to their special flavour, nutritive value and medicinal properties. Mushrooms are a good source of vitamin B, C and D, including niacin, riboflavin, thiamine, and various minerals including potassium, phosphorus, calcium, magnesium, iron and copper. Mushroom is considered to be a nutritious food, rich in protein, low in fat and carbohydrates. However, mushroom growing can help in a long way in the efficient utilization of agricultural and industrial waste. It can also play a significant role to alleviate poverty and generate employment opportunity for educated

unemployed youth. Mushroom farming is increasingly becoming attractive to small farmers because the farmers in rural areas are convinced about the profit of mushroom farming within short time. This can enhance their empowerment to gain other farming skills; greater initial independence and also self-respect. The vocational trainings were given by the Krishi Vigyan Kendra for the farmers, farm-women and unemployed youth to increase the income and make them self-dependent entrepreneur in future. So, the present study was undertaken to find out the knowledge and adoption level of farm women in mushroom cultivation techniques. Mushroom farming is taken up as vocational training programme, as it is considered to provide security to rural poor by providing additional income, thus alleviating the poverty in rural areas and confronting educated unemployment in urban and semi urban areas. It is a subsidiary low cost occupation grown as an indoor crop, by utilizing vertical space and obtaining more productivity per unit area.

Mushrooms are a fleshy fungi having a stem, cap, gills underneath the cap .They can be edible ,wild and toxic too. They provide carbohydrates, but are low in fats and fibers.

MATERIALS AND METHODS

The research methodology for this study involved selecting a sample of farmers who were involved in mushroom cultivation in Nalanda, Bihar. The population for the study consisted of 470 farmers and an updated list of these farmers was prepared by the researcher with the help of the mushroom cultivation center and the KVK. From the population of 470 farmers, a sample of 120 farmers was selected using a random sampling method. In case some of the sampled farmers were not available or did not agree to participate, a reserve list of 10 farmers was prepared to use as a replacement. The data collection was done from the selected farmers to assess their level of knowledge and adoption of mushroom production technology. The questionnaire was designed to collect

information on demographic characteristics, level of knowledge, and adoption of mushroom production technology, as well as the impact of KVK training programs on their knowledge and adoption. The collected data were analyzed using descriptive statistics, such as frequency, percentage, mean, and standard deviation. The study used statistical software to analyze the data and draw conclusions based on the findings.

RESULTS AND DISCUSSION

Age:

The data of Table 1 show that out of 120 respondents, the majority (56.7 per cent) of them were from young age group, 29.2 per cent belong to middle age group, whereas only 14.2 per cent belonged to old age group.

Table - 1 : Age wise distribution of the respondents:

S.N.	Age (Years)	Frequency	Percentage
1.	Young (below 37)	68	56.7
2.	Middle (37-50 years)	35	29.2
3.	Old (above 50)	17	14.2
	Total	120	100.00

From the above findings, it is concluded that majority of the respondents were from young age group, having experienced ups and downs in the life for a considerable period. The findings of the present study were in accordance with the findings of Kaur and Gill (2001).

Education:

The data of Table 2 reveal that out of 120 respondents, the maximum (31.7 per cent) of them were Graduate, 30.00 per cent belonged to Intermediate, 18.3 per cent belonged to Secondary education, 13.3 per cent were Illiterate and 6.7 per cent belonged to Primary education.

Table - 2 : Education wise distribution of the respondents

S. N.	Education	Frequency	Percentage
1.	Illiterate	16	13.3
2.	Primary	8	6.7
3.	Secondary	22	18.3
4.	Intermediate	36	30.0
5.	Graduate	38	31.7
	Total	120	100.00

Highest number of the respondents were graduate i.e 31.7 per cent. The findings of the present study were in accordance with the findings of Noor et al. (2015).

Gender :

The observations with regard to Gender of the respondents are shown in Table 3

Table - 3 : Gender wise distribution of the respondents

S.N.	Gender	Frequency	Percentage
1.	Male	62	51.7
2.	Female	58	48.3
	Total	120	100.00

The data presented in Table 3 revealed that out of 120 respondents the majority (51.7 per cent) was Male and 48.3 per cent of the respondents were Female. Therefore, Most of the respondents were Male.

Marital Status :

The observations with regard to marital status of the respondents are shown in Table 4 .

Table - 4 : Marital status wise distribution of the respondents

S.N.	Marital status	Frequency	Percentage
1.	Married	46	38.3
2.	Unmarried	62	51.7
3.	Widow	12	10.0
	Total	120	100.00

The data presented in Table 4 reveal that out of 120 respondents the majority (51.7 per cent) were unmarried , 38.3 per cent of the respondents were married and in widow category there were 10.0 per cent. Since, most of the respondents were Unmarried.

Caste Category:

It is the social status or position of the respondents which is acquired by heredity. It was categorized into four categories as per Govt. of India norm. The distribution of the respondents according to their categories has been presented in table 5.

Table - 5 : Distribution of Respondents According to Their Caste Category

S.N.	Category	Frequency	Percentage
1.	General	19	15.8
2.	Other Backward class (OBC)	80	66.7
3.	Schedule Caste (SC)	21	17.5
4.	Schedule Tribes(ST)	0	0
	Total	120	100.00

A glance on the Table 5 revealed that maximum (66.7 per cent) number of respondents belonged to the Other Backward Caste followed by Schedule caste 17.5 per cent and General caste 15.8 per cent. No Schedule Tribe participates in the training. This table reported that maximum respondents had Other Backward Caste category. The findings of the present study were in accordance with the findings of Prakash (2000).

Nature of Family:

It was measured based on the information obtained from the respondents whether they are living with more than a pair or with single pair of couples and unmarried children.

This variable was categorized in two categories: joint family and nuclear family.

Table - 6 : Distribution of Respondents According to Their Family Type

S.N.	Nature of Family	Frequency	Percentage
1.	Joint	67	55.8
2.	Nuclear	53	44.2
	Total	120	100.00

The data presented in Table 6 reveal that majority (55.8 per cent) of respondents were from joint family followed by 44.2 per cent respondents from nuclear family. Therefore, it may be concluded that still people in Bihar have faith in joint family. The findings of the present study were in accordance with the findings of Singh *et al.* (2016).

This trend is still prevalent in many parts of the country. The findings of the present study were in accordance with the findings of Singh et al. (2016). They having also reported that 60 per cent of farmers were from joint family.

Agricultural Occupation:

The data presented in Table 7 revealed that 67.5 per cent of the respondents were engaged in agriculture as their main occupation for their livelihood, followed by 26.7 per cent of them were engaged in Secondary occupation, 5.8 per cent of them were engaged in Tertiary occupation.

Table - 7 : Distribution of Respondents According to their Occupation

S.N.	Occupation	Frequency	Percentage
1.	Primary	81	67.5
2.	Secondary	32	26.7
3.	Tertiary	7	5.8
	Total	120	100.00

Knowledge Level of the respondents:

The data presented in Table 8 revealed that majority of the respondents of 63.3 per cent belonged to medium level of knowledge of

mushroom growing followed by 21.7 per cent of the respondents belonged to high level of knowledge and 15 per cent of the respondents were under low level of knowledge.

Table - 8 : Distribution of respondents according to their Knowledge level

S.N.	Knowledge Level	Frequency	Percentage
1.	Low (below 18)	18	15.0
2.	Medium (Between 18-28)	76	63.3
3.	High (Above 28)	26	21.7
Total		120	100.00

Thus, it was concluded that the maximum respondents were belonging to medium level of knowledge.

Adoption Level of the respondents:

The data presented in Table 9 revealed that majority of the respondents of 53.3 per cent belonged to medium level of adoption followed by 24.2 per cent of the respondents who belonged to low level of Adoption and 22.5 per cent of the respondents who were under high level of Adoption.

Thus, it was concluded that the maximum respondents were belonging to medium level of Adoption.

Table - 9 : Distribution of respondents according to their level of adoption

S.N.	Adoption Level	Frequency	Percentage
1.	Low (below 18)	29	24.2
2.	Medium (Between 18-28)	64	53.3
3.	High (Above 28)	27	22.5
Total		120	100.00

CONCLUSION

It was concluded that the majority of farmers involved in mushroom cultivation in District Nalanda, Bihar, were young, male, unmarried, and belonged to the OBC category. Most of them had a graduate education, were engaged in primary occupations and had joint family types. The study found that the majority of respondents had a medium level of knowledge (63.3%) and adoption

(53.3%) of mushroom production technology. This indicates that there is a need for increased awareness and training programs to improve farmers' knowledge and adoption of new technologies in mushroom cultivation.

The findings of the study suggest that KVK training programs have a positive impact on improving the knowledge and adoption of mushroom production technology among farmers. Therefore, there is a need for continued and expanded KVK training programs to enhance the productivity and income of farmers in the agricultural sector, Pant, Hemlata (2010-11).

REFERENCES

1. Kaur, B. and Gill O. K. (2001). Trainees profile of specialized training course in rabbit farming. *Indian J. Extn.Edu.*, 25 : 106-109.

2. Noor M. I; Sanaullah, N; Muhammad Y. S; Mmuhammad U. A; Riaz H. J; Aamir A. M.,Akbar K. K; Sajid A. ,; Rizwan J. and Abdul H. J. (2015). Economics Analysis of Mango Orchard Production under Contract Farming in TalukaTando Adam Districtb Sanghar Sindh. *Pakistan Journal of Biology, Agriculture and Healthcare*, 5(11).

3. Prakash, O; Mahipal and Kherde, R.L. (2000). A study of perception of training needs of landless farm women in scientific dairy farming Practices. *Advance in Agricultural Research in India*, 4:196-209.

4. Singh, S; Sachan, R. and Singh, A.P.(2016). Knowledge and adoption of wheat growers. *Journal of Communication Studies*, Vol. XXXIV, No.1.

5. Pant, Hemlata (2010-11):Mushroom Tatha Mushroom spawn utpadan vidhi, Button, oyster Mushroom Key Swad ki Tulana/Uganey hetu Ichhchit Mushroom Parikshan & Santushti ke sandarbh me interview dwara sarvekchan : Ek addhyan, *Grameen Vikas Sandesh* 5 (1,2) Page No. 46

EFFECCT OF DIFFERENT VARIABLES ON PHYSICO-CHEMICAL STUDY OF TULSI ENRICHED MANGO (MANGIFERA INDICA L.) ICE-CREAM

Siddhartha Singh*, Akhilesh Kumar Singh, Manoj Kumar, Manvendra Singh, Rajesh Kumar Pal and Kautuk Upadhyay

Department of Animal Husbandry and Dairying

T. D. P.G. College, Jaunpur, (U.P.), India

Corresponding Email : akhileshbhu88@gmail.com

Received : 02.11.2022

ABSTRACT

Accepted : 05.12.2022

An experiment was conducted to improve physico-chemical properties of ofTulsi enriched mango Ice-cream. Ice cream with, 2, to 18 % Mango pulp, 0.5 Tulsi extract was prepared. Mango flavor ice cream (without Mango pulp) was kept as control treatment. Ice cream samples were analyzed for physico-chemical parameter such as overrun, protien, fat, meltdown, moisture, pH. The highest score were awarded to the ice cream with 10% mango pulp, 0.5% tulsi extract and 12% sugar followed by ice cream with 18 mango pulp, 0.5% Tulsi extract and 12% sugar. The ice cream samples without mango pulp were liked least. Overrun, protien, fat, meltdown, moisture, pH was affected significantly at 10% mango pulp, 0.5% Tulsi Extract and 12% sugar.

Keywords : Ice cream, mango (*mangifera indica*), pulp physico-chemical analysis, tulsi extract

INTRODUCTION

Ice cream is a frozen dairy product obtained from cow or buffalo milk or a combination thereof from cream and other milk product, with or without addition of cane sugar, egg, fruits juice, preserved fruits, nuts, chocolate, edible flavours and permitted food colours. It may contain added emulsifiers and stabilizers not exceeding 0.5 per cent by weight and should contain minimum of 3.5 per cent protein, 10.00 percent fat, 36.00 percent total solids'. (FSSAI 2017)

Ice cream is a liquid mixture that turns into a paste after simultaneously shaking and cooling although the definition of ice cream varies from country to country due to differing regulations and traditions of composition(Fiol.*et al*, 2016).

Ice cream is a nutritionally enriched frozen dairy product consumed by all groups of age, particularly children, during summer (Sharif *et al.*, 2005). In the ice cream mix that will become ice cream are so many elements of different nature as sugars, fats, dairy, stabilizer, water, among others. And they all have to be correctly blended and emulsified together so there is nothing left behind that may reduce the quality of the final product. Making this possible considering the characteristics and behaviors of each ingredient and the relationships between them is what is known as the balancing exercise. We can make it stable and spreadable at negative temperature from -11 to -18 °C (standard ice cream serving temperatures) (Corvitto, 2011).

Ice-cream is palatable, nutrition, healthful and relatively inexpensive food. One serving good average composition of vanilla ice-cream (1\6 qt) supplies approximately 200 calories, 3.9 g protein, 0.31g calcium, 0.104 g phosphorus , 0.14 g iron, 548 IU vitamin A, 0.038 mg thiamine and 0.236 mg riboflavin. (Arbuckle and Shumaker, 1956)

Indian ice cream industry is one of the fastest growing segments of the dairy or food processing industry. India has a low per capita ice cream consumption of 400 ml as compared with per capital consumption of ice cream of 22 L in the United States and 3 L in China. With the improving cold chain infrastructure in the country coupled with increasing disposable income and the changing lifestyle, the sector has great potential for growth (Anonymous 2018 b).

The ice cream industry in India generated revenue of about USD 1.5 billion in 2016 and is projected to generate revenue of approximately USD 3.4 billion by 2021. Lately, frozen desserts which are made out of vegetable oils have been eating into the market share of ice cream. Key players offering frozen desserts in India are Kwality Walls, Vadilal, and Cream Bell. (Anonymous 2018 b). The annual growth rate of Ice cream is 10-15 per cent. (Singh *et al*, 2003)

Ice cream is highly palatable and nutritious food. One serving of a good Ice cream at average composition (100 g) supplies approximately 200 calories, 4 g protein, 0.13 g calcium, 0.105 g of phosphorous, 0.1 mg iron, 490 IU Vit A, 0.38 mg thiamine and 0.24 mg riboflavin (Bhandari, 2001).

In addition to nutritional importance ice cream can be used as a functional food. The increased awareness of the consumer regarding health and nutrition related issues as well as the role of several food regulator bodies to promote the production and consumption of minimally processed, healthier and more nutritious food products, appear to be steering a transformation within the food industry (Fogliano and Vitaglione, 2005). Dairy products, such as ice creams and

yogurts, are considered nutritious foods and present a great potential to incorporate bio- actives (Aboulfazli *et al.*, 2016). Many research support that ice cream is a good source of different essential nutrients. In related to health, ice cream is used as probiotics carrier and also for other disease prevention.

Tulsi or basil (*Ocimum sanctum.*) has been utilized in Ayurveda, a Hindu form of medicine, for thousands of years for its various medicinal effects. The Charaka Samhita, an ancient Ayurvedic text mentions it. Tulsi or basil is an adaptogen, which means it helps you adapt to stress by harmonizing numerous processes in our body. It is regarded as a kind of "elixir of life" in Ayurveda, with a powerful aroma and astringent taste, and is thought to prolong lifespan. Tulsior basil is pungent and bitter in taste pungent in the post digestive effect and hashot potency. It alleviates kapha and vatadoshas, but slightly aggravates the pitta dosha. It possesses light and dry attributes.

On the contrary the seeds are oily and slimy in attributes and have a cold potency. Tulsi or basil is a stimulant, aromatic herb and effectively reduces the fever. Tulsi or basil also contains antioxidants like beta-carotene that help in preventing cell damage. Tulsi or the holy basil is famous throughout the globe for its healing and other medicinal properties. Its leaves are helpful in sharpening memory and incurring fever and common cold. They also act as an anti-stress agent and also help in purifying blood. This, in turn, helps in reducing the risk of heart attacks and also lowers the cholesterol level. The leaves of the basil are also effective in reducing mouth ulcer and other infections of the mouth. Tulsi belongs to division- Magnoliophyta, genus *Ocimum*, species-*Sanctum*, class Magnoliopsida, order Lamiales and family Lamiaceae. In Hindi and Gujarati, tulsi is known as "Tulsi, in Sanskrit language it is known as 'Tulasi' and in Marathi, it is called as "Tulas' while in English, tulsi is known as "Holy basil" or "Sacred basil". (Krishna *et al.*, 2014)

Tulsi or basil (*Ocimum sanctum* L.) family-Lamiaceae is a fragrant plant that has long been used to treat headaches, coughing, diarrhoea, constipation, warts, worms, and kidney issues. It has a long history as a culinary herb, with its leaves lending a distinct flavour to a wide range of recipes. It also contains insecticidal, nematicidal, fungistatic, and antibacterial biologically active components, as well as aroma compounds and essential oils. Extracts of the tulsi or basil are widely used to treat a variety of ailments, including the common cold, inflammation, malaria, heart disease, headaches, stomach problems, kidney stones, cardiac problems, and more. Tulsi, Indian basil, is also useful for air cleaning. The tulsi plant is an excellent repellent for flies Mosquitoes and other insects. Also, extremely useful in the fight against malaria. (David, 2015)

In sweet basil, the fat content and calorific value is low while high amount of minerals and vitamin A are present. In 2.5 g of basil leaves (five fresh leaves), there are 96.6 IU vitamin A, 3.85 mg calcium, less than 1 calorie, 11.55 mg potassium, and smaller proportions of vitamin C and other vitamins, protein, fibre and minerals. The GRAS (generally recognized as safe) list of the US Department of Agriculture includes sweet basil leaf to be used in the range of 2–680 ppm and 0.01–50 ppm for the essential oil. The use of exceedingly large quantities of oil is suggested to have a health risk due to the occurrence of carcinogenic compounds. The GRAS-suggested amount of basil essential oil is very minute, and internal use of a large amount of this oil should be avoided (Hanifet al., 2011; Hosseini-Parvaret al., 2015).

Mango (*Mangifera indica* L.) belongs to family Anacardiaceae, genus *Mangifera* and species *indica*. It is one of the important tropical fruits and ranks 5th in total production of major fruit crops in the world, after musa, citrus, grapes and apples (Mukherjee, 1997). It is valued for its excellent flavor, attractive fragrance, delicious taste and nutritional benefits. Therefore, it is referred to as the

'King of fruits', in Asia. More than a thousand varieties of mango are grown in the world. However, sensory properties and biochemical constituents vary depending on the mango variety. India produces about 50% of the total world mango fruit production. It was reported that during the 2012-2013, India produced about 16.2 million metric tonnes of mangoes (NHB, 2013). Mango cultivation in the world was reported to be around 3.7 mn ha (Jahurule et al., 2015) and its cultivation in India was estimated to be approximately 2.46 mn ha, which is the highest among the mango growing countries (Sekhar, 2013). Although the production of mango is initially limited to Asian countries, currently its production has spread to different parts of the world such as South and Central America, Africa, Australia, and some parts of Europe especially Spain (Mukeherjee, 1997).

Mango (*Mangifera indica* L.) is one of the favoured fruits in the tropical and subtropical regions. It is often called 'King of Fruits'. It has an excellent flavour, attractive fragrance, delicious taste and high nutritional value that have made it one of the best fruits (Pal, 1998). India accounts almost half of the world production. It is receiving increased attention because of its potential antioxidant activity. It is a rich source of Vitamin A and C. It contains on an average 54 µg of Vitamin A and 36.4 mg Vitamin C per 100 g of edible portion. It also contains minerals like calcium, iron and phosphorous. Carotenoids are compounds of great dietary importance not only as precursors of vitamin A, but also as molecules that take part in cell protection and consumer attraction due to the visual colour they provide to food (Jadhav et al., 2009).

MATERIALS AND METHODS

Raw materials including milk, fresh cream, skim milk powder, sugar, Mango Flavour, and fresh Mango were purchased from local market of Jaunpur Uttar Pradesh.

Table - 1 : Basic steps involved in the ice-cream manufacture as

Sr. No	Process steps	Processing conditions
1.	Selection of the ingredients	Raw material processed and packed under set norms should be of acceptable quality and from approved suppliers.
2.	Weighing of the ingredients.	-
3.	Mixing of the ingredients	Sequence of addition depends on type of ingredients used.
4.	Pasteurization of the mix.	By one of three methods: a) Batch or long hold at 68.3°C(155°F)/30 min b) High temperature short time 79.4°C (175°F)/25 s c) Ultra high temperature at 150°C (300°F) for a fraction of a second.
5.	Homogenization of the mix.	At 70-75°C by two stages. (I stage 2500 psi and II stage 500 psi)
6.	Cooling of the mix	To 4°C (40°F) or below
7.	Ageing of the mix	At 2-4°C/2-24 h
8.	Freezing and aeration	At a high shear / 20 -30 s extrusion temperature -5 to -6°C.
9.	Packaging	In case of low temperature continuous freezer -9°C or below. At the extrusion temperature or after hardening in the blast tunnel.
10.	Hardening of the ice-cream	At -25 to -30°C
11.	Storage of the ice-cream	At -12.2 to -23°C (10 to -10°F)

Source: - Khanna (2004) suggested the steps for a typical manufacturing process of ice-cream as under Investigation.

Different treatments used in this Investigation

T_0 = Ice-cream mix as per standard

T_1 = Ice-cream mix as per standard + Acceptable level of 0.5% Tulsi extract + 2% Mango pulp

T_2 = Ice-cream mix as per standard + Acceptable level of 0.5% Tulsi extract + 4% Mango pulp

T_3 = Ice-cream mix as per standard + Acceptable level of 0.5% Tulsi extract + 6% Mango pulp

T_4 = Ice-cream mix as per standard + Acceptable level of 0.5% Tulsi extract + 8% Mango pulp

T_5 = Ice-cream mix as per standard + Acceptable level of 0.5% Tulsi extract + 10% Mango

pulp

T_6 = Ice-cream mix as per standard + Acceptable level of 0.5% Tulsi extract + 12% Mango pulp

T_7 = Ice-cream mix as per standard + Acceptable level of 0.5% Tulsi extract + 14% Mango pulp



T_8 = Ice-cream mix as per standard + Acceptable level of 0.5% Tulsi extract + 16% Mango pulp

T_9 = Ice-cream mix as per standard + Acceptable level of 0.5% Tulsi extract + 18% Mango pulp

For ice cream preparation, all the ingredients were weighed using weighing balance according to the formulation. All the dry ingredients as well as liquid ingredients were manually mixed thoroughly by stirring until a uniform mixture resulted. The ice cream mix was pasteurized at 72 °C for 30 minutes as described by Marshall and Arbuckl (1996) to destroy pathogenic organisms. After pasteurization, the mix was homogenized in electric homogenizer as described by Berger and White (1976). The fat globules were reduced in size by homogenization in order to obtain a uniform dispersion of the fat. After homogenization, mango pulp, Tulsi extract was incorporated according to treatment.

The mix was cooled down to 4 °C immediately in a deep freezer and stored at 4 °C for ageing. Ageing also increased the viscosity of the mix. After ageing, the mix was frozen at -5 °C in the freezing chamber of electrically operated batch type ice cream machine. When the desired consistency had been attained, the product was filled into disposable paper cups. The cups were immediately transferred to the hardening unit maintained at -250C to -300C and the ice cream were kept for 24 hours. These cups were then transferred to deep freezer at -20 °C. The data obtained in the research on different aspects were tabulated and analysed statically using the methods of One-way analysis of variance (ANOVA).

Procedure of Methodology-



RESULTS AND DISCUSSION

Results of chemical analysis are shown in Table 2. On the basis of physico chemical parameter such as pH, moisture, fat, protein and Overrun percentage the ice cream with 0.5% Tulsi Extract + 10% mango pulp has shown significant as compared to other treatment. Significant decrease in pH and significant increase in acidity is due to presence of ascorbic acid in mangos. The results of this study are

supported by the findings of(Akhilesh et al 2018)

Table - 2 : Physico-chemical properties of mango pulp enriched ice cream

Treatment	Moisture	Overrun percentage	Fat	pH	Protein
T0	64.63	42.08	11.70	6.35	4.13
T1	64.78	41.32	11.88	6.32	4.18
T2	65.45	40.63	11.78	6.29	4.13
T3	65.94	40.20	11.82	6.33	3.98
T4	66.33	38.49	11.70	6.31	4.23
T5	65.13	36.85	11.74	6.29	4.26
T6	65.87	36.85	11.44	6.28	4.08
T7	66.35	40.65	11.37	6.30	3.90
T8	64.75	37.87	11.19	6.28	4.20
T9	65.99	36.20	11.14	6.28	4.06

Table 2 Shows that the Moisture score was the highest in treatment T₇(66.35%) than the lowest score was recorded in treatment T₀ (64.63%). The Protein score of prepared Ice cream was highest in treatment T₅(4.26%) and the lowest in treatment T₃ (3.98%). The maximum and minimum score for Fat was recorded in treatment T₁(11.88%) the lowest in treatment T₉(11.14%). The pH score was highest in treatment T₀(6.35) and the lowest value recorded in treatment in T₆ (6.28),T₈ (6.28)and T₉ (6.28). The Overrun percentage was highest in treatment T₁ (42.08%) and the lowest value recorded in treatment in T₆(36.85%) and T₇(36.85%). It is clear from table 1 that the score obtained by treatment T1 for sensory parameters as color& appearance, flavor, body & texture, melting properties and overall acceptability was (8.73, 8.37, 8.75, 8.11, 8.41 respectively) highest whereas the score obtained by Control sample (T0) for color& appearance flavor, body & texture,melting properties and overall acceptability was (7.81, 7.66, 7.9, 7.16, 6.69 respectively) very near to the treatment T1. It is revealed from the above result that the difference in the value obtained by treatment T1 was Non - Significant (p<0.05) to the value obtained by T0 (Control) treatment.The results of this study are supported by the findings of Gwyszczynska and Kaluziak (1971).

CONCLUSION

A new variety of ice creams particularly rich in chewy eating sensation and pleasant flavor can be manufactured by utilizing frozen strawberry fruit.

Appearance, taste and mouthfeel characteristics of the ice cream improved upon the addition of the fruit. Further research is recommended to establish the conditions for processing of strawberry fruit in different forms and levels of addition in ice cream without compromising the quality of the product. On the basis of above findings it can be concluded that the superior quality of Tulsi extract enriched Mango Ice Cream can be prepared by addition of 10% of mango pulp, 0.5% tulsi Extract and 10% of sugar as the overall acceptance for treatment combination T1 was highest in all physico-chemical treatment.

REFERENCES

1. Aboulfazli, F., Shori, A. B., & Baba, A. S., (2016). Effects of the replacement of cow milk with vegetable milk on probiotics and nutritional profile of fermented ice cream. *LWT - Food Science and Technology*, 70(1), 261-270. <http://dx.doi.org/10.1016/j.lwt.2016.02.056>.
2. Akhilesh Kumar Singh, DC Rai, Uday Pratap Singh and Sudhir Kumar (2018). Effect of different variables on physico-chemical properties of Ashwagandha enriched strawberry pulp ice cream. *The Pharma Innovation Journal* 2018; 7(4): 440-443
3. Anonymous, b. 2018. A study of india's ice-cream market 2018.
4. Arbuckle, W.S. and Shumaker, M.L. (1956). Milk solids not fat in ice cream. *Dairy Sci. Abstr.*, 12:236
5. Berger, K. G., and G. W. White. 1976. The fat globule membrane in ice cream. *Dairy Ind. Int.* 41:199
6. Bhandari, V. (2001) Ice-cream manufacture and technology. The textbook published by Tata Mc.Graw.Hill publishing company limited, New Delhi.
7. Corvito, A., (2011). The secrets of ice cream lose. *Del helado ice cream without Secrets*. Sant Cugat Del Valles, Vilbo. Spain.
8. David, J. 2015. Preparation of herbal shrikhand prepared with basil (*Ocimum basilicum*) extract. *The Pharma Innovation Journal*, 4(8): 81-84.
9. Fiol C., Prado, D., Romero. Laburu, N., Mora, M., and Alawa, J.I., (2017). Introduction of a new family of ice creams. *International Journal of Gastronomy and Food Science*,
10. Fogliano V, Vitaglione P., (2005). Functional foods: planning and development. *Mol Nutr Food Res.* 49:256-62
11. FSSAI, 2006. Food Safety and Standards Act, 2006 (www.fssai.gov.in)
12. Gwyszczynska, A. and H. Kaluziak, 1971. Changes in ice cream during storage. *Prezmysl Spozywczy*, 25: 66-9 (*Food Sci. Tech. Abstr.*, 18: 68; 1986)
13. Hanif, M.A., Al-Maskari, M.Y., Al-Maskari, A., Al-Shukaili, A., Al-Maskari, A.Y. and Al-Sabahi, J.N. (2011) Essential oil composition, antimicrobial and antioxidant activities of unexplored Omani basil. *Journal of Medicinal Plants Research* 5, 751-757.
14. Hosseini-Parvar, S.H., Matia-Merino, L. and Golding, M. (2015) Effect of basil seed gum (BSG) on textural, rheological and microstructural properties of model processed cheese. *Food Hydrocolloids* 44, 557-567.
15. Jadhav, P. V., Kawadkar, D. K., Kshirsagar, R. B., Bansode, V. V. and Jadhao, A. S., 2009. Studies on development of carotene rich mango powder. *Ind J. of Nutri Diet*, 46 (3): 112-117.
16. Jahurul MHA, Norulaini NAN, Zaidul ISM, Jinap S, Sahena F, Azmir J, Sharif KM, Mohd Omar AK. Cocoa butter fats and possibilities of substitution in food products concerning cocoa varieties, alternative

60	<i>Siddhartha Singh et. al.</i>	
<p>sources, extraction methods, composition, and characteristics. J Food Engineer 2013, 117(4):467–476.</p> <p>17. Krishna S. G., Bhavani, R. T. and Kumar, P. 2014. Tulsi- the wonder herh (Pharmacological Activities of Ocimum sanctum). American Journal of Ethnomedicine.1 (1):089-095.</p> <p>18. Marshall, R.T. and Arbuckle, W.S. (1996) Ice Cream. 5th Edition, Chapman & Hall, New York. http://dx.doi.org/10.1007/978-1-4613-0477-7</p> <p>19. Mukherjee SK. Introduction: Botany and importance. In: R.E. Litz (Ed.). The mango: Botanay, production and uses. 1997:1-19, CRC Press Cab International, USA.</p> <p>20. National Horticulture Board (NHB). Ministry of agriculture. Gurgaon (Haryana) India: Government of India 2013.</p>	<p>21. Pal, R. K., 1998. Ripening and rheological properties of mango as influenced by ethereal and carbide. J. Food Technol., 35 (4): 358-360.</p> <p>22. Sekhar C, Selvarajan M, Pounraj A, Prahadeeswaran M. Production and export of mango in India-A paradigm to the developing nations. Am Intern Res Human Arts Social Sci 2013, 4:78–87.</p> <p>23. Sharif, N., K.S. Ghanghesh, Y.A.S. Gnan and A. Rahouma, (2005). Bacteriological quality of ice cream in Tripoli-Libya. Food Control, 17: 637-641.</p> <p>24. Singh, P.K., J. Singh and R.K. Pandey, 2003.Consumer acceptance of Soft serve ice cream. Nutritional seminar on value added dairy products Dec. 18 (2005).</p>	